

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification<sup>7</sup> : <b>C12N 9/00</b></p>	<p><b>A2-</b></p>	<p>(11) International Publication Number: <b>WO 00/60059</b> (43) International Publication Date: 12 October 2000 (12.10.00)</p>
<p>(21) International Application Number: PCT/DK00/00148 (22) International Filing Date: 28 March 2000 (28.03.00) (30) Priority Data: PA 1999 00437 30 March 1999 (30.03.99) DK (71) Applicant: NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK). (72) Inventors: ANDERSEN, Carsten; Højeloft Vænge 162, DK-3500 Værløse (DK). JØRGENSEN, Christel, Thea; Livjærggade 41, 4.th., DK-2100 Copenhagen Ø (DK). BISGÅRD-FRANTZEN, Henrik; Elmevænget 8B, DK-2880 Bagsværd (DK). SVENDSEN, Allan; Bakkeledet 28, DK-3460 Birkerød (DK). KJÆRULFF, Søren; Kongsdalsvej 47, DK-2720 Vanløse (DK).</p>		<p>(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: ALPHA-AMYLASE VARIANTS</p> <p>(57) Abstract</p> <p>The invention relates to a variant of a parent Termamyl-like alpha-amylase, which variant exhibits altered properties, in particular reduced capability of cleaving a substrate close to the branching point, and improved substrate specificity and/or improved specific activity relative to the parent alpha-amylase.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## Alpha-AMYLASE VARIANTS

### FIELD OF THE INVENTION

The present invention relates, *inter alia*, to novel variants  
5 of parent Termamyl-like alpha-amylases, notably variants  
exhibiting altered properties, in particular altered cleavage  
pattern (relative to the parent) which are advantageous with  
respect to applications of the variants in, in particular,  
industrial starch processing (e.g., starch liquefaction or  
10 saccharification).

### BACKGROUND OF THE INVENTION

Alpha-Amylases (alpha-1,4-glucan-4-glucanohydrolases, EC  
3.2.1.1) constitute a group of enzymes which catalyze hydrolysis  
15 of starch and other linear and branched 1,4-glucosidic oligo-  
and polysaccharides.

There is a very extensive body of patent and scientific  
literature relating to this industrially very important class of  
enzymes. A number of alpha-amylase such as Termamyl-like  
20 alpha-amylases variants are known from, e.g., WO 90/11352, WO  
95/10603, WO 95/26397, WO 96/23873, WO 96/23874 and WO 97/41213.

Among recent disclosure relating to alpha-amylases, WO  
96/23874 provides three-dimensional, X-ray crystal structural  
data for a Termamyl-like alpha-amylase, referred to as BA2,  
25 which consists of the 300 N-terminal amino acid residues of the  
*B. amyloliquefaciens* alpha-amylase comprising the amino acid  
sequence shown in SEQ ID NO: 6 herein and amino acids 301-483 of  
the C-terminal end of the *B. licheniformis* alpha-amylase  
comprising the amino acid sequence shown in SEQ ID NO: 4 herein  
30 (the latter being available commercially under the tradename  
Termamyl™), and which is thus closely related to the  
industrially important *Bacillus* alpha-amylases (which in the  
present context are embraced within the meaning of the term  
"Termamyl-like alpha-amylases", and which include, *inter alia*,  
35 the *B. licheniformis*, *B. amyloliquefaciens* and *B.*  
*stearothermophilus* alpha-amylases). WO 96/23874 further  
describes methodology for designing, on the basis of an analysis

of the structure of a parent Termamyl-like alpha-amylase, variants of the parent Termamyl-like alpha-amylase which exhibit altered properties relative to the parent.

WO 96/23874 and WO 97/41213 (Novo Nordisk) discloses  
5 Termamyl-like alpha-amylase variants with an altered cleavage pattern containing mutations in the amino acid residues V54, D53, Y56, Q333, G57 and A52 of the sequence shown in SEQ ID NO: 4 herein.

#### 10 BRIEF DISCLOSURE OF THE INVENTION

The present invention relates to novel alpha-amylolytic variants (mutants) of a Termamyl-like alpha-amylase, in particular variants exhibiting altered cleavage pattern (relative to the parent), which are advantageous in connection  
15 with the industrial processing of starch (starch liquefaction, saccharification and the like).

The inventors have surprisingly found variants with altered properties, in particular altered cleavage pattern which have improved reduced capability of cleaving an substrate close to  
20 the branching point, and further have improved substrate specificity and/or improved specific activity, in comparison to the WO 96/23874 and WO 97/41213 (Novo Nordisk) disclosed Termamyl-like alpha-amylase variants with an altered cleavage pattern containing mutations in the amino acid residues V54,  
25 D53, Y56, Q333, G57 and A52 of the sequence shown in SEQ ID NO: 4 herein.

The invention further relates to DNA constructs encoding variants of the invention, to composition comprising variants of the invention, to methods for preparing variants of the  
30 invention, and to the use of variants and compositions of the invention, alone or in combination with other alpha-amylolytic enzymes, in various industrial processes, e.g., starch liquefaction, and in detergent compositions, such as laundry, dish washing and hard surface cleaning compositions; ethanol  
35 production, such as fuel, drinking and industrial ethanol production; desizing of textiles, fabrics or garments etc.

### Nomenclature

In the present description and claims, the conventional one-letter and three-letter codes for amino acid residues are used.

For ease of reference, alpha-amylase variants of the invention are described by use of the following nomenclature:

Original amino acid(s):position(s):substituted amino acid(s)

According to this nomenclature, for instance the substitution of alanine for asparagine in position 30 is shown as:

10 Ala30Asn or A30N

a deletion of alanine in the same position is shown as:

Ala30\* or A30\*

and insertion of an additional amino acid residue, such as lysine, is shown as:

15 \*30aLys or \*30aK

A deletion of a consecutive stretch of amino acid residues, such as amino acid residues 30-33, is indicated as (30-33)\* or  $\Delta$ (A30-N33) or delta(A30-N33).

Where a specific alpha-amylase contains a "deletion" in comparison with other alpha-amylases and an insertion is made in such a position this is indicated as:

\*36aAsp or \*36aD

for insertion of an aspartic acid in position 36

Multiple mutations are separated by plus signs, i.e.:

25 Ala30Asp + Glu34Ser or A30N+E34S

representing mutations in positions 30 and 34 substituting alanine and glutamic acid for asparagine and serine, respectively. Multiple mutations may also be separated as follows, i.e., meaning the same as the plus sign:

30 Ala30Asp/Glu34Ser or A30N/E34S

When one or more alternative amino acid residues may be inserted in a given position it is indicated as

A30N,E or

A30N or A30E

35 Furthermore, when a position suitable for modification is identified herein without any specific modification being suggested, or A30X, it is to be understood that any amino acid

residue may be substituted for the amino acid residue present in the position. Thus, for instance, when a modification of an alanine in position 30 is mentioned, but not specified, or specified as "X", it is to be understood that the alanine may be  
5 deleted or substituted for any other amino acid, i.e., any one of: R,N,D,C,Q,E,G,H,I,L,K,M,F,P,S,T,W,Y,V.

#### DETAILED DISCLOSURE OF THE INVENTION

##### The Termamyl-like alpha-amylase

10 It is well known that a number of alpha-amylases produced by *Bacillus* spp. are highly homologous on the amino acid level. For instance, the *B. licheniformis* alpha-amylase comprising the amino acid sequence shown in SEQ ID NO: 4 (commercially available as Termamyl™) has been found to be about 89%  
15 homologous with the *B. amyloliquefaciens* alpha-amylase comprising the amino acid sequence shown in SEQ ID NO: 6 and about 79% homologous with the *B. stearothermophilus* alpha-amylase comprising the amino acid sequence shown in SEQ ID NO: 8. Further homologous alpha-amylases include an alpha-amylase  
20 derived from a strain of the *Bacillus* sp. NCIB 12289, NCIB 12512, NCIB 12513 or DSM 9375, all of which are described in detail in WO 95/26397, and the #707 alpha-amylase described by Tsukamoto et al., Biochemical and Biophysical Research Communications, 151 (1988), pp. 25-31.

25 Still further homologous alpha-amylases include the alpha-amylase produced by the *B. licheniformis* strain described in EP 0252666 (ATCC 27811), and the alpha-amylases identified in WO 91/00353 and WO 94/18314. Other commercial Termamyl-like *B. licheniformis* alpha-amylases are Optitherm™ and Takatherm™  
30 (available from Solvay), Maxamyl™ (available from Gist-brocades/Genencor), Spezym AA™ and Spezyme Delta AA™ (available from Genencor), and Keistase™ (available from Daiwa).

Because of the substantial homology found between these alpha-amylases, they are considered to belong to the same class  
35 of alpha-amylases, namely the class of "Termamyl-like alpha-amylases".

Accordingly, in the present context, the term "Termamyl-like alpha-amylase" is intended to indicate an alpha-amylase, which at the amino acid level exhibits a substantial homology to Termamyl<sup>TM</sup>, i.e., the *B. licheniformis* alpha-amylase having the amino acid sequence shown in SEQ ID NO: 4 herein. In other words, a Termamyl-like alpha-amylase is an alpha-amylase, which has the amino acid sequence shown in SEQ ID NO: 2, 4, 6, or 8 herein, and the amino acid sequence shown in SEQ ID NO: 1 or 2 of WO 95/26397 or in Tsukamoto et al., 1988, or i) which displays at least 60%, preferred at least 70%, more preferred at least 75%, even more preferred at least 80%, especially at least 85%, especially preferred at least 90%, even especially more preferred at least 95% homology, more preferred at least 97%, more preferred at least 99% with at least one of said amino acid sequences and/or ii) displays immunological cross-reactivity with an antibody raised against at least one of said alpha-amylases, and/or iii) is encoded by a DNA sequence which hybridises to the DNA sequences encoding the above-specified alpha-amylases which are apparent from SEQ ID NOS: 1, 3, 5 and 7 of the present application and SEQ ID NOS: 4 and 5 of WO 95/26397, respectively.

In connection with property i), the "homology" may be determined by use of any conventional algorithm, preferably by use of the GAP programme from the GCG package version 7.3 (June 1993) using default values for GAP penalties, which is a GAP creation penalty of 3.0 and GAP extension penalty of 0.1, (Genetic Computer Group (1991) Programme Manual for the GCG Package, version 7, 575 Science Drive, Madison, Wisconsin, USA 53711).

A structural alignment between Termamyl and a Termamyl-like alpha-amylase may be used to identify equivalent/corresponding positions in other Termamyl-like alpha-amylases. One method of obtaining said structural alignment is to use the Pile Up programme from the GCG package using default values of gap penalties, i.e., a gap creation penalty of 3.0 and gap extension penalty of 0.1. Other structural alignment methods include the hydrophobic cluster analysis (Gaboriaud et al.,

(1987), FEBS LETTERS 224, pp. 149-155) and reverse threading (Huber, T ; Torda, AE, PROTEIN SCIENCE Vol. 7, No. 1 pp. 142-149 (1998). Property ii) of the alpha-amylase, i.e., the immunological cross reactivity, may be assayed using an antibody raised against, or reactive with, at least one epitope of the relevant Termamyl-like alpha-amylase. The antibody, which may either be monoclonal or polyclonal, may be produced by methods known in the art, e.g., as described by Hudson et al., Practical Immunology, Third edition (1989), Blackwell Scientific Publications. The immunological cross-reactivity may be determined using assays known in the art, examples of which are Western Blotting or radial immunodiffusion assay, e.g., as described by Hudson et al., 1989. In this respect, immunological cross-reactivity between the alpha-amylases having the amino acid sequences SEQ ID NOS: 2, 4, 6, or 8, respectively, have been found.

The oligonucleotide probe used in the characterization of the Termamyl-like alpha-amylase in accordance with property iii) above may suitably be prepared on the basis of the full or partial nucleotide or amino acid sequence of the alpha-amylase in question.

Suitable conditions for testing hybridization involve presoaking in 5xSSC and prehybridizing for 1 hour at -40°C in a solution of 20% formamide, 5xDenhardt's solution, 50mM sodium phosphate, pH 6.8, and 50mg of denatured sonicated calf thymus DNA, followed by hybridization in the same solution supplemented with 100mM ATP for 18 hours at -40°C, followed by three times washing of the filter in 2xSSC, 0.2% SDS at 40°C for 30 minutes (low stringency), preferred at 50°C (medium stringency), more preferably at 65°C (high stringency), even more preferably at ~75°C (very high stringency). More details about the hybridization method can be found in Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor, 1989.

In the present context, "derived from" is intended not only to indicate an alpha-amylase produced or producible by a strain of the organism in question, but also an alpha-amylase encoded



by a DNA sequence isolated from such strain and produced in a host organism transformed with said DNA sequence. Finally, the term is intended to indicate an alpha-amylase, which is encoded by a DNA sequence of synthetic and/or cDNA origin and which has the identifying characteristics of the alpha-amylase in question. The term is also intended to indicate that the parent alpha-amylase may be a variant of a naturally occurring alpha-amylase, i.e. a variant, which is the result of a modification (insertion, substitution, deletion) of one or more amino acid residues of the naturally occurring alpha-amylase.

#### Parent hybrid alpha-amylases

The parent alpha-amylase may be a hybrid alpha-amylase, i.e., an alpha-amylase, which comprises a combination of partial amino acid sequences derived from at least two alpha-amylases.

The parent hybrid alpha-amylase may be one, which on the basis of amino acid homology and/or immunological cross-reactivity and/or DNA hybridization (as defined above) can be determined to belong to the Termamyl-like alpha-amylase family. In this case, the hybrid alpha-amylase is typically composed of at least one part of a Termamyl-like alpha-amylase and part(s) of one or more other alpha-amylases selected from Termamyl-like alpha-amylases or non-Termamyl-like alpha-amylases of microbial (bacterial or fungal) and/or mammalian origin.

Thus, the parent hybrid alpha-amylase may comprise a combination of partial amino acid sequences deriving from at least two Termamyl-like alpha-amylases, or from at least one Termamyl-like and at least one non-Termamyl-like bacterial alpha-amylase, or from at least one Termamyl-like and at least one fungal alpha-amylase. The Termamyl-like alpha-amylase from which a partial amino acid sequence derives may, e.g., be any of those specific Termamyl-like alpha-amylases referred to herein.

For instance, the parent alpha-amylase may comprise a C-terminal part of an alpha-amylase derived from a strain of *B. licheniformis*, and a N-terminal part of an alpha-amylase derived from a strain of *B. amyloliquefaciens* or from a strain of *B. stearothermophilus*. For instance, the parent alpha-amylase may

comprise at least 430 amino acid residues of the C-terminal part of the *B. licheniformis* alpha-amylase, and may, e.g., comprise a) an amino acid segment corresponding to the 37 N-terminal amino acid residues of the *B. amyloliquefaciens* alpha-amylase having the amino acid sequence shown in SEQ ID NO: 6 and an amino acid segment corresponding to the 445 C-terminal amino acid residues of the *B. licheniformis* alpha-amylase having the amino acid sequence shown in SEQ ID NO: 4, or b) an amino acid segment corresponding to the 68 N-terminal amino acid residues of the *B. stearothermophilus* alpha-amylase having the amino acid sequence shown in SEQ ID NO: 8 and an amino acid segment corresponding to the 415 C-terminal amino acid residues of the *B. licheniformis* alpha-amylase having the amino acid sequence shown in SEQ ID NO: 4.

In a preferred embodiment the parent Termamyl-like alpha-amylase is a hybrid Termamyl-like alpha-amylase identical to the *Bacillus licheniformis* alpha-amylase shown in SEQ ID NO: 4, except that the N-terminal 35 amino acid residues (of the mature protein) is replaced with the N-terminal 33 amino acid residues of the mature protein of the *Bacillus amyloliquefaciens* alpha-amylase (BAN) shown in SEQ ID NO: 6. Said hybrid may further have the following mutations: H156Y+A181T+N190F+A209V+Q264S (using the numbering in SEQ ID NO: 4) referred to as LE174.

Another preferred parent hybrid alpha-amylase is LE429 shown in SEQ ID NO: 2.

The non-Termamyl-like alpha-amylase may, e.g., be a fungal alpha-amylase, a mammalian or a plant alpha-amylase or a bacterial alpha-amylase (different from a Termamyl-like alpha-amylase). Specific examples of such alpha-amylases include the *Aspergillus oryzae* TAKA alpha-amylase, the *A. niger* acid alpha-amylase, the *Bacillus subtilis* alpha-amylase, the porcine pancreatic alpha-amylase and a barley alpha-amylase. All of these alpha-amylases have elucidated structures, which are markedly different from the structure of a typical Termamyl-like alpha-amylase as referred to herein.

The fungal alpha-amylases mentioned above, i.e., derived from *A. niger* and *A. oryzae*, are highly homologous on the amino

acid level and generally considered to belong to the same family of alpha-amylases. The fungal alpha-amylase derived from *Aspergillus oryzae* is commercially available under the tradename Fungamyl™.

5 Furthermore, when a particular variant of a Termamyl-like alpha-amylase (variant of the invention) is referred to - in a conventional manner - by reference to modification (e.g., deletion or substitution) of specific amino acid residues in the amino acid sequence of a specific Termamyl-like alpha-amylase,  
10 it is to be understood that variants of another Termamyl-like alpha-amylase modified in the equivalent position(s) (as determined from the best possible amino acid sequence alignment between the respective amino acid sequences) are encompassed thereby.

15 A preferred embodiment of a variant of the invention is one derived from a *B. licheniformis* alpha-amylase (as parent Termamyl-like alpha-amylase), e.g., one of those referred to above, such as the *B. licheniformis* alpha-amylase having the amino acid sequence shown in SEQ ID NO: 4.

20

#### Construction of variants of the invention

The construction of the variant of interest may be accomplished by cultivating a microorganism comprising a DNA sequence encoding the variant under conditions which are  
25 conducive for producing the variant. The variant may then subsequently be recovered from the resulting culture broth. This is described in detail further below.

#### Altered properties

30 The following discusses the relationship between mutations, which may be present in variants of the invention, and desirable alterations in properties (relative to those of a parent Termamyl-like alpha-amylase), which may result there from.

In the first aspect the invention relates to a variant of  
35 a parent Termamyl-like alpha-amylase, comprising an alteration at one or more positions selected from the group of:

W13, G48, T49, S50, Q51, A52, D53, V54, G57, G107, G108, A111,

S168, M197, wherein (a) the alteration(s) are independently

(i) an insertion of an amino acid downstream of the amino acid which occupies the position,

(ii) a deletion of the amino acid which occupies the position, or

(iii) a substitution of the amino acid which occupies the position with a different amino acid,

(b) the variant has alpha-amylase activity and (c) each position corresponds to a position of the amino acid sequence of the parent Termamyl-like alpha-amylase having the amino acid sequence of SEQ ID NO: 4.

In a preferred embodiment the above variants of the invention comprise a mutation in a position corresponding to at least one of the following mutations in the amino acid sequence shown in SEQ ID NO: 4:

V54N, A52S, A52S+V54N, T49L, T49+G107A, A52S+V54N+T49L+G107A, A52S+V54N+T49L, G107A, Q51R, Q51R+A52S, A52N; or  
T49F+G107A, T49V+G107A, T49D+G107A, T49Y+G107A, T49S+G107A, T49N+G107A, T49I+G107A, T49L+A52S+G107A, T49L+A52T+G107A, T49L+A52F+G107A, T49L+A52L+G107A, T49L+A52I+G107A, T49L+A52V+G107A; or  
T49V, T49I, T49D, T49N, T49S, T49Y, T49F, T49W, T49M, T49E, T49Q, T49K, T49R, A52T, A52L, A52I, A52V, A52M, A52F, A52Y, A52W, V54M, G107V, G07I, G107L, G107C.

In a preferred embodiment a variant of the invention comprises at least one mutation in a position corresponding to the following mutations in the amino acid sequence shown in SEQ ID NO: 4:

W13F,L,I,V,Y,A;  
G48A,V,S,T,I,L;  
\*48aD or \*48aY (i.e., insertion of D or Y);  
T49X;  
\*49aX (i.e., insertion of any possible amino acid residue)  
S50X, in particular D,Y,L,T,V,I;  
Q51R,K;  
A52X, in particular A52S,N,T,F,L,I,V;  
D53E,Q,Y,I,N,S,T,V,L;

V54X, in particular V54I,N,W,Y,F,L;  
G57S,A,V,L,I,F,Y,T;  
G107X, in particular G107A,V,S,T,I,L,C;  
G108X, in particular G108A,V,S,T,I,L;  
5 A111V,I,L;  
S168Y;  
M197X, in particular Y,F,L,I,T,A,G.

In a preferred embodiment a variant of the invention comprises the following mutations corresponding to the following  
10 mutations in the amino acid sequence shown in SEQ ID NO: 4:  
T49X+A52X+V54N/I/L/Y/F/W+G107A, and may further comprise G108A.

In a preferred embodiment a variant of the invention comprises at least one mutation corresponding to the following mutations in the amino acid sequence shown in SEQ ID NO: 4:

15 T49L+G107A;  
T49I+G107A;  
T49L+G107A+V54I;  
T49I+G107A+V54I;  
A52S+V54N+T49L+G107A;  
20 A52S+V54I+T49L+G107A;  
A52S+T49L+G107A;  
A52T+T49L+G107A;  
A52S+V54N+T49I+G107A;  
A52S+V54I+T49I+G107A;  
25 A52S+T49I+G107A;  
T49L+G108A;  
T49I+G108A;  
T49L+G108A+V54I;  
T49I+G108A+V54I.

30 All of the above-mentioned variants of the invention have altered properties (meaning increased or decreased properties), in particular at least one of the following properties relative to the parent alpha-amylase: reduced ability to cleave a substrate close to the branching point, improved substrate  
35 specificity and/or improved specific activity, altered substrate binding, altered thermal stability, altered pH/activity profile, altered pH/stability profile, altered stability towards

oxidation, altered  $\text{Ca}^{2+}$  dependency.

5    Stability

        In the context of the present invention, mutations (including amino acid substitutions and/or deletions) of importance with respect to achieving altered stability, in particular improved stability (i.e., higher or lower), at  
10 especially low pH (i.e., pH 4-6) include any of the mutations listed in the in "Altered properties" section, above and the variants mentioned right below.

        The following variants: Q360A,K; N102A, N326A,L, N190G, N190K; Y262A,K,E (using the BAN, i.e., SEQ ID N: 6, numbering)  
15 were also tested for pH stability. A preferred parent alpha-amylase may be BA2 described above. The pH stability was determined as described in the "Materials & Methods" section.

20     $\text{Ca}^{2+}$  stability

        Altered  $\text{Ca}^{2+}$  stability means the stability of the enzyme under  $\text{Ca}^{2+}$  depletion has been improved, i.e., higher or lower stability. In the context of the present invention, mutations (including amino acid substitutions) of importance with respect  
25 to achieving altered  $\text{Ca}^{2+}$  stability, in particular improved  $\text{Ca}^{2+}$  stability, i.e., higher or lower stability, at especially low pH (i.e., pH 4-6) include any of the mutations listed in the in "Altered properties" section above.

30    Specific activity

        In a further aspect of the present invention, important mutations with respect to obtaining variants exhibiting altered specific activity, in particular increased or decreased specific activity, especially at temperatures from 60-100°C, preferably  
35 70-95°C, especially 80-90°C, include any of the mutations listed in the in "Altered properties" section above.

        The specific activity of LE174 and LE429 was determined to

16,000 NU/mg using the Phadebas® assay described in the "Materials and Methods" section.

#### Altered cleavage pattern

5 In the starch liquefaction process it is desirable to use an alpha-amylase, which is capable of degrading the starch molecules into long, branched oligosaccharides, rather than an alpha-amylase, which gives rise to formation of shorter, branched oligosaccharides (like conventional Termamyl-like  
10 alpha-amylases). Short, branched oligosaccharides (panose precursors) are not hydrolyzed satisfactorily by pullulanases, which are used after alpha-amylase treatment in the liquefaction process, or simultaneously with a saccharifying amyloglucosidase (glucoamylase), or before adding a saccharifying  
15 amyloglucosidase (glucoamylase). Thus, in the presence of panose precursors, the product mixture present after the glucoamylase treatment contains a significant proportion of short, branched, so-called limit-dextrin, viz. the trisaccharide panose. The presence of panose lowers the saccharification yield sig-  
20 nificantly and is thus undesirable.

It has been reported previously (US patent 5,234,823) that, when saccharifying with glucoamylase and pullulanase, the presence of residual alpha-amylase activity arising from the liquefaction process, can lead to lower yields of glucose,  
25 if the alpha-amylase is not inactivated before the saccharification stage. This inactivation can be typically carried out by adjusting the pH to below 4.7 at 95°C, before lowering the temperature to 60°C for saccharification.

The reason for this negative effect on glucose yield is  
30 not fully understood, but it is assumed that the liquefying alpha-amylase (for example Termamyl 120 L from *B.licheniformis*) generates "limit dextrins" (which are poor substrates for pullulanase), by hydrolysing 1,4-alpha-glucosidic linkages close to and on both sides of the  
35 branching points in amylopectin. Hydrolysis of these limit dextrins by glucoamylase leads to a build up of the trisaccharide panose, which is only slowly hydrolysed by

glucoamylase.

The development of a thermostable alpha-amylase, which does not suffer from this disadvantage, would be a significant improvement, as no separate inactivation step would be required.

Thus, the aim of the present invention is to arrive at a mutant alpha-amylase having appropriately modified starch-degradation characteristics but retaining the thermostability of the parent Termamyl-like alpha-amylase.

Accordingly, the invention relates to a variant of a Termamyl-like alpha-amylase, which has an improved reduced ability to cleave a substrate close to the branching point, and further has improved substrate specificity and/or improved specific activity.

Of particular interest is a variant, which cleaves an amylopectin substrate, from the reducing end, more than one glucose unit from the branching point, preferably more than two or three glucose units from the branching point, i.e., at a further distance from the branching point than that obtained by use of a wild type *B. licheniformis* alpha-amylase.

It may be mentioned here that according to WO 96/23874, variants comprising at least one of the following mutations are expected to prevent cleavage close to the branching point:

V54L, I, F, Y, W, R, K, H, E, Q;

D53L, I, F, Y, W;

Y56W;

Q333W;

G57, all possible amino acid residues;

A52, amino acid residues larger than A, e.g., A52W, Y, L, F, I.

Mutations of particular interest in relation to obtaining variants according to the invention having an improved reduced ability to cleave a substrate close to the branching point, and further has improved substrate specificity and/or improved specific activity include mutations at the following positions in *B. licheniformis* alpha-amylase, SEQ ID NO: 4:

H156, A181, N190, A209, Q264 and I201.

It should be emphasised that not only the Termamyl-like



alpha-amylases mentioned specifically below may be used. Also other commercial Termamyl-like alpha-amylases can be used. An unexhaustive list of such alpha-amylases is the following:

Alpha-amylases produced by the *B. licheniformis* strain described  
5 in EP 0252666 (ATCC 27811), and the alpha-amylases identified in  
WO 91/00353 and WO 94/18314. Other commercial Termamyl-like *B.*  
*licheniformis* alpha-amylases are Optitherm™ and Takatherm™  
(available from Solvay), Maxamyl™ (available from Gist-  
brocades/Genencor), Spezym AA™ Spezyme Delta AA™ (available  
10 from Genencor), and Keistase™ (available from Daiwa).

All Termamyl-like alpha-amylase may suitably be used as backbone for preparing variants of the invention.

In a preferred embodiment of the invention the parent Termamyl-like alpha-amylase is a hybrid alpha-amylase of SEQ ID  
15 NO: 4 and SEQ ID NO: 6. Specifically, the parent hybrid Termamyl-like alpha-amylase may be a hybrid alpha-amylase comprising the 445 C-terminal amino acid residues of the *B. licheniformis* alpha-amylase shown in SEQ ID NO: 4 and the 37 N-terminal amino acid residues of the mature alpha-amylase derived  
20 from *B. amyloliquefaciens* shown in SEQ ID NO: 6, which may suitably further have the following mutations: H156Y+A181T+N190F+A209V+Q264S (using the numbering in SEQ ID NO: 4). This hybrid is referred to as LE174. The LE174 hybrid may be combined with a further mutation I201F to form a parent hybrid  
25 Termamyl-like alpha-amylase having the following mutations H156Y+A181T+N190F+A209V+Q264S+I201F (using SEQ ID NO: 4 for the numbering). This hybrid variant is shown in SEQ ID NO: 2 and is used in the examples below, and is referred to as LE429.

Also, LE174 or LE429 (SEQ ID NO: 2) or *B. licheniformis*  
30 alpha-amylase shown in SEQ ID NO: 4 comprising one or more of the following mutations may be used as backbone (using SEQ ID NO: 4 for the numbering of the mutations):

E119C;

S130C;

35 D124C;

R127C;

A52all possible amino acid residues;  
 S85all possible amino acid residues;  
 N96all possible amino acid residues;  
 V129all possible amino acid residues;  
 5 A269all possible amino acid residues;  
 A378all possible amino acid residues;  
 S148all possible amino acid residues, in particular S148N;  
 E211all possible amino acid residues, in particular E211Q;  
 N188all possible amino acid residues, in particular N188S, N188P  
 10 M197all possible amino acid residues, in particular M197T,  
 M197A, M197G, M197I, M197L, M197Y, M197F, M197I;  
 W138all possible amino acid residues, in particular W138Y;  
 D207all possible amino acid residues, in particular D207Y;  
 H133all possible amino acid residues, in particular H133Y;  
 15 H205all possible amino acid residues, in particular H205H,  
 H205C, H205R;  
 S187all possible amino acid residues, in particular S187D;  
 A210all possible amino acid residues, in particular A210S,  
 A210T;  
 20 H405all possible amino acid residues, in particular H405D;  
 K176all possible amino acid residues, in particular K176R;  
 F279all possible amino acid residues, in particular F279Y;  
 Q298all possible amino acid residues, in particular Q298H;  
 G299all possible amino acid residues, in particular G299R;  
 25 L308all possible amino acid residues, in particular L308F;  
 T412all possible amino acid residues, in particular T412A;

Further, *B. licheniformis* alpha-amylase shown in SEQ ID NO:  
 4 comprising at least one of the following mutations may be used  
 as backbone:

30 M15all possible amino acid residues;  
 A33all possible amino acid residues;

When using LE429 (shown in SEQ ID NO: 2) as the backbone  
 (i.e., as the parent Termamyl-like alpha-amylase) by combining  
 LE174 with the mutation I201F (SEQ ID NO: 4 numbering), the  
 35 mutations/alterations, in particular substitutions, deletions  
 and insertions, may according to the invention be made in one or  
 more of the following positions to improve the reduced ability

to cleave a substrate close to the branching point, and to improve substrate specificity and/or improved specific activity: W13, G48, T49, S50, Q51, A52, D53, V54, G57, G107, G108, A111, S168, M197 (using the SEQ ID NO: 4 numbering)

5 wherein (a) the alteration(s) are independently

(i) an insertion of an amino acid downstream of the amino acid which occupies the position,

(ii) a deletion of the amino acid which occupies the position, or

10 (iii) a substitution of the amino acid which occupies the position with a different amino acid,

(b) the variant has alpha-amylase activity and (c) each position corresponds to a position of the amino acid sequence of the parent Termamyl-like alpha-amylase having the amino acid sequence of SEQ ID NO: 4.

15 In a preferred embodiment a variant of the invention comprises at least one mutation in a position corresponding to the following mutations in the amino acid sequence shown in SEQ ID NO: 4:

20 V54N, A52S, A52S+V54N, T49L, T49+G107A, A52S+V54N+T49L+G107A, A52S+V54N+T49L, G107A, Q51R, Q51R+A52S, A52N; or  
T49F+G107A, T49V+G107A, T49D+G107A, T49Y+G107A, T49S+G107A, T49N+G107A, T49I+G107A, T49L+A52S+G107A, T49L+A52T+G107A, T49L+A52F+G107A, T49L+A52L+G107A, T49L+A52I+G107A,  
25 T49L+A52V+G107A; or  
T49V, T49I, T49D, T49N, T49S, T49Y, T49F, T49W, T49M, T49E, T49Q, T49K, T49R, A52T, A52L, A52I, A52V, A52M, A52F, A52Y, A52W, V54M, G107V, G07I, G107L, G107C.

30 In a preferred embodiment a variant of the invention comprises at least one mutation in a position corresponding to the following mutations in the amino acid sequence shown in SEQ ID NO: 4:

W13F, L, I, V, Y, A;

G48A, V, S, T, I, L;

35 \*48aD or \*48aY (i.e., insertion of D or Y);

T49X;

\*49aX (i.e., insertion of any amino acid residue)

S50X, in particular D,Y,L,T,V,I;  
Q51R,K;  
A52X, in particular A52S,N,T,F,L,I,V;  
D53E,Q,Y,I,N,S,T,V,L;  
5 V54X, in particular V54I,N,W,Y,F,L;  
G57S,A,V,L,I,F,Y,T;  
G107X, in particular G107A,V,S,T,I,L,C;  
G108X, in particular G108A,V,S,T,I,L;  
A111V,I,L;  
10 S168Y;  
M197X, in particular Y,F,L,I,T,A,G.

In a preferred embodiment a variant of the invention comprises at least one mutation in a position corresponding to the following mutations in the amino acid sequence shown in SEQ  
15 ID NO: 4:  
T49X+A52X+V54N/I/L/Y/F/W+G107A, and may further comprise G108A.

In a preferred embodiment a variant of the invention comprises at least one mutation in a position corresponding to the following mutations in the amino acid sequence shown in SEQ  
20 ID NO: 4:  
T49L+G107A;  
T49I+G107A;  
T49L+G107A+V54I;  
T49I+G107A+V54I;  
25 A52S+V54N+T49L+G107A;  
A52S+V54I+T49L+G107A;  
A52S+T49L+G107A;  
A52T+T49L+G107A;  
A52S+V54N+T49I+G107A;  
30 A52S+V54I+T49I+G107A;  
A52S+T49I+G107A;  
T49L+G108A;  
T49I+G108A;  
T49L+G108A+V54I;  
35 T49I+G108A+V54I.

General mutations in variants of the invention

It may be preferred that a variant of the invention comprises one or more modifications in addition to those outlined above. Thus, it may be advantageous that one or more proline residues present in the part of the alpha-amylase variant which is modified is/are replaced with a non-proline residue which may be any of the possible, naturally occurring non-proline residues, and which preferably is an alanine, glycine, serine, threonine, valine or leucine.

Analogously, it may be preferred that one or more cysteine residues present among the amino acid residues with which the parent alpha-amylase is modified is/are replaced with a non-cysteine residue such as serine, alanine, threonine, glycine, valine or leucine.

Furthermore, a variant of the invention may - either as the only modification or in combination with any of the above outlined modifications - be modified so that one or more Asp and/or Glu present in an amino acid fragment corresponding to the amino acid fragment 185-209 of SEQ ID NO. 4 is replaced by an Asn and/or Gln, respectively. Also of interest is the replacement, in the Termamyl-like alpha-amylase, of one or more of the Lys residues present in an amino acid fragment corresponding to the amino acid fragment 185-209 of SEQ ID NO: 4 by an Arg.

It will be understood that the present invention encompasses variants incorporating two or more of the above outlined modifications.

Furthermore, it may be advantageous to introduce point-mutations in any of the variants described herein.

#### Methods for preparing alpha-amylase variants

Several methods for introducing mutations into genes are known in the art. After a brief discussion of the cloning of alpha-amylase-encoding DNA sequences, methods for generating mutations at specific sites within the alpha-amylase-encoding sequence will be discussed.

Cloning a DNA sequence encoding an alpha-amylase

The DNA sequence encoding a parent alpha-amylase may be isolated from any cell or microorganism producing the alpha-amylase in question, using various methods well known in the art. First, a genomic DNA and/or cDNA library should be constructed using chromosomal DNA or messenger RNA from the organism that produces the alpha-amylase to be studied. Then, if the amino acid sequence of the alpha-amylase is known, homologous, labelled oligonucleotide probes may be synthesized and used to identify alpha-amylase-encoding clones from a genomic library prepared from the organism in question. Alternatively, a labelled oligonucleotide probe containing sequences homologous to a known alpha-amylase gene could be used as a probe to identify alpha-amylase-encoding clones, using hybridization and washing conditions of lower stringency.

Yet another method for identifying alpha-amylase-encoding clones would involve inserting fragments of genomic DNA into an expression vector, such as a plasmid, transforming alpha-amylase-negative bacteria with the resulting genomic DNA library, and then plating the transformed bacteria onto agar containing a substrate for alpha-amylase, thereby allowing clones expressing the alpha-amylase to be identified.

Alternatively, the DNA sequence encoding the enzyme may be prepared synthetically by established standard methods, e.g., the phosphoroamidite method described by S.L. Beaucage and M.H. Caruthers (1981) or the method described by Matthes et al. (1984). In the phosphoroamidite method, oligonucleotides are synthesized, e.g., in an automatic DNA synthesizer, purified, annealed, ligated and cloned in appropriate vectors.

Finally, the DNA sequence may be of mixed genomic and synthetic origin, mixed synthetic and cDNA origin or mixed genomic and cDNA origin, prepared by ligating fragments of synthetic, genomic or cDNA origin (as appropriate, the fragments corresponding to various parts of the entire DNA sequence), in accordance with standard techniques. The DNA sequence may also be prepared by polymerase chain reaction (PCR) using specific primers, for instance as described in US 4,683,202 or R.K. Saiki

et al. (1988).

#### Site-directed mutagenesis

Once an alpha-amylase-encoding DNA sequence has been  
5 isolated, and desirable sites for mutation identified, mutations  
may be introduced using synthetic oligonucleotides. These oligo-  
nucleotides contain nucleotide sequences flanking the desired  
mutation sites; mutant nucleotides are inserted during oligo-  
nucleotide synthesis. In a specific method, a single-stranded  
10 gap of DNA, bridging the alpha-amylase-encoding sequence, is  
created in a vector carrying the alpha-amylase gene. Then the  
synthetic nucleotide, bearing the desired mutation, is annealed  
to a homologous portion of the single-stranded DNA. The re-  
maining gap is then filled in with DNA polymerase I (Klenow  
15 fragment) and the construct is ligated using T4 ligase. A  
specific example of this method is described in Morinaga et al.  
(1984). US 4,760,025 disclose the introduction of oligonucleoti-  
des encoding multiple mutations by performing minor alterations  
of the cassette. However, an even greater variety of mutations  
20 can be introduced at any one time by the Morinaga method,  
because a multitude of oligonucleotides, of various lengths, can  
be introduced.

Another method for introducing mutations into alpha-amylase-  
encoding DNA sequences is described in Nelson and Long (1989).  
25 It involves the 3-step generation of a PCR fragment containing  
the desired mutation introduced by using a chemically syn-  
thesized DNA strand as one of the primers in the PCR reactions.  
From the PCR-generated fragment, a DNA fragment carrying the  
mutation may be isolated by cleavage with restriction  
30 endonucleases and reinserted into an expression plasmid.

#### Random Mutagenesis

Random mutagenesis is suitably performed either as localised  
or region-specific random mutagenesis in at least three parts of  
35 the gene translating to the amino acid sequence shown in  
question, or within the whole gene.

The random mutagenesis of a DNA sequence encoding a parent

alpha-amylase may be conveniently performed by use of any method known in the art.

In relation to the above, a further aspect of the present invention relates to a method for generating a variant of a parent alpha-amylase, e.g., wherein the variant exhibits a reduced capability of cleaving an oligo-saccharide substrate close to the branching point, and further exhibits improved substrate specificity and/or improved specific activity relative to the parent, the method:

- 10 (a) subjecting a DNA sequence encoding the parent alpha-amylase to random mutagenesis,
- (b) expressing the mutated DNA sequence obtained in step (a) in a host cell, and
- (c) screening for host cells expressing an alpha-amylase
- 15 variant which has an altered property (i.e., thermal stability) relative to the parent alpha-amylase.

Step (a) of the above method of the invention is preferably performed using doped primers. For instance, the random mutagenesis may be performed by use of a suitable physical or chemical mutagenizing agent, by use of a suitable oligonucleotide, or by subjecting the DNA sequence to PCR generated mutagenesis. Furthermore, the random mutagenesis may be performed by use of any combination of these mutagenizing agents. The mutagenizing agent may, e.g., be one, which induces

20 transitions, transversions, inversions, scrambling, deletions, and/or insertions.

Examples of a physical or chemical mutagenizing agent suitable for the present purpose include ultraviolet (UV) ir-radiation, hydroxylamine, N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), O-methyl hydroxylamine, nitrous acid, ethyl methane sulphonate (EMS), sodium bisulphite, formic acid, and nucleotide analogues.

30 When such agents are used, the mutagenesis is typically performed by incubating the DNA sequence encoding the parent enzyme to be mutagenized in the presence of the mutagenizing agent of choice under suitable conditions for the mutagenesis to take place, and selecting for mutated DNA having the desired properties. When the mutagenesis is performed by the use of an



oligonucleotide, the oligonucleotide may be doped or spiked with the three non-parent nucleotides during the synthesis of the oligonucleotide at the positions, which are to be changed. The doping or spiking may be done so that codons for unwanted amino acids are avoided. The doped or spiked oligonucleotide can be incorporated into the DNA encoding the alpha-amylase enzyme by any published technique, using e.g., PCR, LCR or any DNA polymerase and ligase as deemed appropriate. Preferably, the doping is carried out using "constant random doping", in which the percentage of wild type and mutation in each position is predefined. Furthermore, the doping may be directed toward a preference for the introduction of certain nucleotides, and thereby a preference for the introduction of one or more specific amino acid residues. The doping may be made, e.g., so as to allow for the introduction of 90% wild type and 10% mutations in each position. An additional consideration in the choice of a doping scheme is based on genetic as well as protein-structural constraints. The doping scheme may be made by using the DOPE program, which, inter alia, ensures that introduction of stop codons is avoided. When PCR-generated mutagenesis is used, either a chemically treated or non-treated gene encoding a parent alpha-amylase is subjected to PCR under conditions that increase the mis-incorporation of nucleotides (Deshler 1992; Leung et al., Technique, Vol.1, 1989, pp. 11-15). A mutator strain of *E. coli* (Fowler et al., Molec. Gen. Genet., 133, 1974, pp. 179-191), *S. cerevisiae* or any other microbial organism may be used for the random mutagenesis of the DNA encoding the alpha-amylase by, e.g., transforming a plasmid containing the parent glycosylase into the mutator strain, growing the mutator strain with the plasmid and isolating the mutated plasmid from the mutator strain. The mutated plasmid may be subsequently transformed into the expression organism. The DNA sequence to be mutagenized may be conveniently present in a genomic or cDNA library prepared from an organism expressing the parent alpha-amylase. Alternatively, the DNA sequence may be present on a suitable vector such as a plasmid or a bacteriophage, which as such may be incubated with or

otherwise exposed to the mutagenising agent. The DNA to be mutagenized may also be present in a host cell either by being integrated in the genome of said cell or by being present on a vector harboured in the cell. Finally, the DNA to be mutagenized  
5 may be in isolated form. It will be understood that the DNA sequence to be subjected to random mutagenesis is preferably a cDNA or a genomic DNA sequence. In some cases it may be convenient to amplify the mutated DNA sequence prior to performing the expression step b) or the screening step c). Such  
10 amplification may be performed in accordance with methods known in the art, the presently preferred method being PCR-generated amplification using oligonucleotide primers prepared on the basis of the DNA or amino acid sequence of the parent enzyme. Subsequent to the incubation with or exposure to the  
15 mutagenising agent, the mutated DNA is expressed by culturing a suitable host cell carrying the DNA sequence under conditions allowing expression to take place. The host cell used for this purpose may be one which has been transformed with the mutated DNA sequence, optionally present on a vector, or one which was  
20 carried the DNA sequence encoding the parent enzyme during the mutagenesis treatment. Examples of suitable host cells are the following: gram positive bacteria such as *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*, *Bacillus brevis*, *Bacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus*  
25 *amyloliquefaciens*, *Bacillus coagulans*, *Bacillus circulans*, *Bacillus lautus*, *Bacillus megaterium*, *Bacillus thuringiensis*, *Streptomyces lividans* or *Streptomyces murinus*; and gram-negative bacteria such as *E. coli*. The mutated DNA sequence may further comprise a DNA sequence encoding functions permitting expression  
30 of the mutated DNA sequence.

#### Localised random mutagenesis

The random mutagenesis may be advantageously localised to a part of the parent alpha-amylase in question. This may, e.g., be advantageous when certain regions of the enzyme have been identified to be of particular importance for a given property of the enzyme, and when modified are expected to result in a variant having improved properties. Such regions may normally be identified when the tertiary structure of the parent enzyme has been elucidated and related to the function of the enzyme.

The localised, or region-specific, random mutagenesis is conveniently performed by use of PCR generated mutagenesis techniques as described above or any other suitable technique known in the art. Alternatively, the DNA sequence encoding the part of the DNA sequence to be modified may be isolated, e.g., by insertion into a suitable vector, and said part may be subsequently subjected to mutagenesis by use of any of the mutagenesis methods discussed above.

#### Alternative methods of providing alpha-amylase variants

Alternative methods for providing variants of the invention include gene-shuffling method known in the art including the methods e.g., described in WO 95/22625 (from Affymax Technologies N.V.) and WO 96/00343 (from Novo Nordisk A/S).

#### Expression of alpha-amylase variants

According to the invention, a DNA sequence encoding the variant produced by methods described above, or by any alternative methods known in the art, can be expressed, in enzyme form, using an expression vector which typically includes control sequences encoding a promoter, operator, ribosome binding site, translation initiation signal, and, optionally, a repressor gene or various activator genes.

The recombinant expression vector carrying the DNA sequence encoding an alpha-amylase variant of the invention may be any vector, which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the

host cell into which it is to be introduced. Thus, the vector may be an autonomously replicating vector, i.e., a vector, which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid, a bacteriophage or an extrachromosomal element, minichromosome or an artificial chromosome. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

In the vector, the DNA sequence should be operably connected to a suitable promoter sequence. The promoter may be any DNA sequence, which shows transcriptional activity in the host cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell. Examples of suitable promoters for directing the transcription of the DNA sequence encoding an alpha-amylase variant of the invention, especially in a bacterial host, are the promoter of the lac operon of *E.coli*, the *Streptomyces coelicolor* agarase gene *dagA* promoters, the promoters of the *Bacillus licheniformis* alpha-amylase gene (*amyL*), the promoters of the *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), the promoters of the *Bacillus amyloliquefaciens* alpha-amylase (*amyQ*), the promoters of the *Bacillus subtilis* *xylA* and *xylB* genes etc. For transcription in a fungal host, examples of useful promoters are those derived from the gene encoding *A. oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *A. niger* neutral alpha-amylase, *A. niger* acid stable alpha-amylase, *A. niger* glucoamylase, *Rhizomucor miehei* lipase, *A. oryzae* alkaline protease, *A. oryzae* triose phosphate isomerase or *A. nidulans* acetamidase.

The expression vector of the invention may also comprise a suitable transcription terminator and, in eukaryotes, polyadenylation sequences operably connected to the DNA sequence encoding the alpha-amylase variant of the invention. Termination and polyadenylation sequences may suitably be derived from the same sources as the promoter.

The vector may further comprise a DNA sequence enabling the

vector to replicate in the host cell in question. Examples of such sequences are the origins of replication of plasmids pUC19, pACYC177, pUB110, pE194, pAMB1 and pIJ702.

The vector may also comprise a selectable marker, e.g., a gene the product of which complements a defect in the host cell, such as the *dal* genes from *B. subtilis* or *B. licheniformis*, or one which confers antibiotic resistance such as ampicillin, kanamycin, chloramphenicol or tetracyclin resistance. Furthermore, the vector may comprise *Aspergillus* selection markers such as *amdS*, *argB*, *niaD* and *sC*, a marker giving rise to hygromycin resistance, or the selection may be accomplished by co-transformation, e.g., as described in WO 91/17243.

While intracellular expression may be advantageous in some respects, e.g., when using certain bacteria as host cells, it is generally preferred that the expression is extracellular. In general, the *Bacillus* alpha-amylases mentioned herein comprise a pre-region permitting secretion of the expressed protease into the culture medium. If desirable, this pre-region may be replaced by a different preregion or signal sequence, conveniently accomplished by substitution of the DNA sequences encoding the respective preregions.

The procedures used to ligate the DNA construct of the invention encoding an alpha-amylase variant, the promoter, terminator and other elements, respectively, and to insert them into suitable vectors containing the information necessary for replication, are well known to persons skilled in the art (cf., for instance, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor, 1989).

The cell of the invention, either comprising a DNA construct or an expression vector of the invention as defined above, is advantageously used as a host cell in the recombinant production of an alpha-amylase variant of the invention. The cell may be transformed with the DNA construct of the invention encoding the variant, conveniently by integrating the DNA construct (in one or more copies) in the host chromosome. This integration is generally considered to be an advantage as the DNA sequence is more likely to be stably maintained in the cell. Integration of

the DNA constructs into the host chromosome may be performed according to conventional methods, e.g., by homologous or heterologous recombination. Alternatively, the cell may be transformed with an expression vector as described above in connection with the different types of host cells.

The cell of the invention may be a cell of a higher organism such as a mammal or an insect, but is preferably a microbial cell, e.g., a bacterial or a fungal (including yeast) cell.

Examples of suitable bacteria are gram-positive bacteria such as *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*, *Bacillus brevis*, *Bacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*, *Bacillus circulans*, *Bacillus lautus*, *Bacillus megaterium*, *Bacillus thuringiensis*, or *Streptomyces lividans* or *Streptomyces murinus*, or gramnegative bacteria such as *E.coli*. The transformation of the bacteria may, for instance, be effected by protoplast transformation or by using competent cells in a manner known per se.

The yeast organism may favourably be selected from a species of *Saccharomyces* or *Schizosaccharomyces*, e.g., *Saccharomyces cerevisiae*. The filamentous fungus may advantageously belong to a species of *Aspergillus*, e.g., *Aspergillus oryzae* or *Aspergillus niger*. Fungal cells may be transformed by a process involving protoplast formation and transformation of the protoplasts followed by regeneration of the cell wall in a manner known per se. A suitable procedure for transformation of *Aspergillus* host cells is described in EP 238 023.

In yet a further aspect, the present invention relates to a method of producing an alpha-amylase variant of the invention, which method comprises cultivating a host cell as described above under conditions conducive to the production of the variant and recovering the variant from the cells and/or culture medium.

The medium used to cultivate the cells may be any conventional medium suitable for growing the host cell in question and obtaining expression of the alpha-amylase variant of the invention. Suitable media are available from commercial suppliers or

may be prepared according to published recipes (e.g., as described in catalogues of the American Type Culture Collection).

5 The alpha-amylase variant secreted from the host cells may conveniently be recovered from the culture medium by well-known procedures, including separating the cells from the medium by centrifugation or filtration, and precipitating proteinaceous components of the medium by means of a salt such as ammonium sulphate, followed by the use of chromatographic procedures such  
10 as ion exchange chromatography, affinity chromatography, or the like.

#### Industrial applications

The alpha-amylase variants of this invention possess  
15 valuable properties allowing for a variety of industrial applications. In particular, enzyme variants of the invention are applicable as a component in washing, dishwashing and hard surface cleaning detergent compositions. Numerous variants are particularly useful in the production of sweeteners and ethanol,  
20 e.g., fuel, drinking or industrial ethanol, from starch, and/or for textile desizing. Conditions for conventional starch-conversion processes, including starch liquefaction and/or saccharification processes, are described in, e.g., US 3,912,590 and in EP patent publications Nos. 252 730 and 63 909.

25

#### Production of sweeteners from starch:

A "traditional" process for conversion of starch to fructose syrups normally consists of three consecutive enzymatic processes, viz. a liquefaction process followed by a sacchari-  
30 fication process and an isomerization process. During the liquefaction process, starch is degraded to dextrins by an alpha-amylase (e.g., Termamyl™) at pH values between 5.5 and 6.2 and at temperatures of 95-160°C for a period of approx. 2 hours. In order to ensure optimal enzyme stability under these condi-  
35 tions, 1 mM of calcium is added (40 ppm free calcium ions).

After the liquefaction process the dextrins are converted into dextrose by addition of a glucoamylase (e.g., AMG™) and a

debranching enzyme, such as an isoamylase or a pullulanase (e.g., Promozyme™). Before this step the pH is reduced to a value below 4.5, maintaining the high temperature (above 95°C), and the liquefying alpha-amylase activity is denatured. The temperature is lowered to 60°C, and glucoamylase and debranching enzyme are added. The saccharification process proceeds for 24-72 hours.

After the saccharification process the pH is increased to a value in the range of 6-8, preferably pH 7.5, and the calcium is removed by ion exchange. The dextrose syrup is then converted into high fructose syrup using, e.g., an immobilized glucoseisomerase (such as Sweetzyme™).

At least one enzymatic improvement of this process could be envisaged: Reduction of the calcium dependency of the liquefying alpha-amylase. Addition of free calcium is required to ensure adequately high stability of the alpha-amylase, but free calcium strongly inhibits the activity of the glucoseisomerase and needs to be removed, by means of an expensive unit operation, to an extent, which reduces the level of free calcium to below 3-5 ppm. Cost savings could be obtained if such an operation could be avoided and the liquefaction process could be performed without addition of free calcium ions.

To achieve that, a less calcium-dependent Termamyl-like alpha-amylase which is stable and highly active at low concentrations of free calcium (< 40 ppm) is required. Such a Termamyl-like alpha-amylase should have a pH optimum at a pH in the range of 4.5-6.5, preferably in the range of 4.5-5.5.

The invention also relates to a composition comprising a mixture of one or more variants of the invention derived from (as the parent Termamyl-like alpha-amylase) the *B. stearothermophilus* alpha-amylase having the sequence shown in SEQ ID NO: 8 and a Termamyl-like alpha-amylase derived from the *B. licheniformis* alpha-amylase having the sequence shown in SEQ ID NO: 4.

Further, the invention also relates to a composition comprising a mixture of one or more variants according the



invention derived from (as the parent Termamyl-like alpha-amylase) the *B. stearothermophilus* alpha-amylase having the sequence shown in SEQ ID NO: 8 and a hybrid alpha-amylase comprising a part of the *B. amyloliquefaciens* alpha-amylase shown in SEQ ID NO: 6 and a part of the *B. licheniformis* alpha-amylase shown in SEQ ID NO: 4. The latter mentioned hybrid Termamyl-like alpha-amylase comprises the 445 C-terminal amino acid residues of the *B. licheniformis* alpha-amylase shown in SEQ ID NO: 4 and the 37 N-terminal amino acid residues of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6. Said latter mentioned hybrid alpha-amylase may suitably comprise the following mutations: H156Y+A181T+N190F+A209V+Q264S (using the numbering in SEQ ID NO: 4) Preferably, said latter mentioned hybrid alpha-amylase may suitably comprise the following mutations: H156Y+A181T+N190F+A209V+Q264S+I201F (using the SEQ ID NO: 4 numbering). In the examples below said last-mentioned parent hybrid Termamyl-like alpha-amylase referred to as LE429 (shown in SEQ ID NO: 2) is used for preparing variants of the invention, which variants may be used in compositions of the invention.

An alpha-amylase variant of the invention or a composition of the invention may in an aspect of the invention be used for starch liquefaction, in detergent composition, such as laundry, dish wash compositions and hard surface cleaning, ethanol production, such as fuel, drinking and industrial ethanol production, desizing of textile, fabric and garments.

#### MATERIALS AND METHODS

##### Enzymes:

LE174: hybrid alpha-amylase variant:

LE174 is a hybrid Termamyl-like alpha-amylase being identical to the Termamyl sequence, i.e., the *Bacillus licheniformis* alpha-amylase shown in SEQ ID NO: 4, except that the N-terminal 35 amino acid residues (of the mature protein) has been replaced by the N-terminal 33 residues of BAN (mature protein), i.e., the *Bacillus amyloliquefaciens* alpha-amylase shown in SEQ ID NO: 6, which further have following mutations:

H156Y+A181T+N190F+A209V+Q264S (SEQ ID NO: 4).

**LE429 hybrid alpha-amylase variant:**

LE429 is a hybrid Termamyl-like alpha-amylase being identical  
5 to the Termamyl sequence, i.e., the *Bacillus licheniformis*  
alpha-amylase shown in SEQ ID NO: 4, except that the N-  
terminal 35 amino acid residues (of the mature protein) has  
been replaced by the N-terminal 33 residues of BAN (mature  
protein), i.e., the *Bacillus amyloliquefaciens* alpha-amylase  
10 shown in SEQ ID NO: 6, which further have following mutations:  
H156Y+A181T+N190F+A209V+Q264S+I201F (SEQ ID NO: 4). LE429 is  
shown as SEQ ID NO: 2 and was constructed by SOE-PCR (Higuchi  
et al. 1988, Nucleic Acids Research 16:7351).

Dextrozyme™ E: a balanced mixture of glucoamylase (AMG) and  
15 pullulanase obtainable from selected strains of *Aspergillus*  
*niger* and *Bacillus deramificans* (available from Novo Nordisk  
A/S)

Fermentation and purification of alpha-amylase variants

20 A *B. subtilis* strain harbouring the relevant expression  
plasmid is streaked on an LB-agar plate with 10 micro g/ml  
kanamycin from -80°C stock, and grown overnight at 37°C.  
The colonies are transferred to 100 ml BPX media supplemented  
with 10 micro g/ml kanamycin in a 500 ml shaking flask.

25 Composition of BPX medium:

Potato starch	100	g/l
Barley flour	50	g/l
BAN 5000 SKB	0.1	g/l
Sodium caseinate	10	g/l
30 Soy Bean Meal	20	g/l
Na <sub>2</sub> HPO <sub>4</sub> , 12 H <sub>2</sub> O	9	g/l
Pluronic™	0.1	g/l

The culture is shaken at 37°C at 270 rpm for 5 days.

35 Cells and cell debris are removed from the fermentation  
broth by centrifugation at 4500 rpm in 20-25 minutes. Afterwards  
the supernatant is filtered to obtain a completely clear

solution. The filtrate is concentrated and washed on an UF-filter (10000 cut off membrane) and the buffer is changed to 20mM Acetate pH 5.5. The UF-filtrate is applied on a S-sepharose F.F. and elution is carried out by step elution with 0.2M NaCl in the same buffer. The eluate is dialysed against 10mM Tris, pH 9.0 and applied on a Q-sepharose F.F. and eluted with a linear gradient from 0-0.3M NaCl over 6 column volumes. The fractions that contain the activity (measured by the Phadebas assay) are pooled, pH was adjusted to pH 7.5 and remaining color was removed by a treatment with 0.5% W/vol. active coal in 5 minutes.

#### Activity determination - (KNU)

One Kilo alpha-amylase Unit (1 KNU) is the amount of enzyme which breaks down 5.26 g starch (Merck, Amylum Solubile, Erg. B 6, Batch 9947275) per hour in Novo Nordisk's standard method for determination of alpha-amylase based upon the following condition:

Substrate	soluble starch
Calcium content in solvent	0.0043 M
Reaction time	7-20 minutes
Temperature	37°C
pH	5.6

Detailed description of Novo Nordisk's analytical method (AF 9) is available on request.

#### Assay for Alpha-Amylase Activity

Alpha-Amylase activity is determined by a method employing Phadebas® tablets as substrate. Phadebas tablets (Phadebas® Amylase Test, supplied by Pharmacia Diagnostic) contain a cross-linked insoluble blue-coloured starch polymer, which has been mixed with bovine serum albumin and a buffer substance and tabletted.

For every single measurement one tablet is suspended in a tube containing 5 ml 50 mM Britton-Robinson buffer (50 mM acetic acid, 50 mM phosphoric acid, 50 mM boric acid, 0.1 mM CaCl<sub>2</sub>, pH adjusted to the value of interest with NaOH). The test is

performed in a water bath at the temperature of interest. The alpha-amylase to be tested is diluted in x ml of 50 mM Britton-Robinson buffer. 1 ml of this alpha-amylase solution is added to the 5 ml 50 mM Britton-Robinson buffer. The starch is hydrolysed  
5 by the alpha-amylase giving soluble blue fragments. The absorbance of the resulting blue solution, measured spectrophotometrically at 620 nm, is a function of the alpha-amylase activity.

It is important that the measured 620 nm absorbance after 10  
10 or 15 minutes of incubation (testing time) is in the range of 0.2 to 2.0 absorbance units at 620 nm. In this absorbance range there is linearity between activity and absorbance (Lambert-Beer law). The dilution of the enzyme must therefore be adjusted to fit this criterion. Under a specified set of conditions (temp.,  
15 pH, reaction time, buffer conditions) 1 mg of a given alpha-amylase will hydrolyse a certain amount of substrate and a blue colour will be produced. The colour intensity is measured at 620 nm. The measured absorbance is directly proportional to the specific activity (activity/mg of pure alpha-amylase protein) of  
20 the alpha-amylase in question under the given set of conditions.

#### Determining Specific Activity

The specific activity is determined using the Phadebas assay (Pharmacia) as activity/mg enzyme.

25

#### Measuring the pH activity profile (pH stability)

The variant is stored in 20 mM TRIS pH 7.5, 0.1 mM,  $\text{CaCl}_2$ , and tested at 30°C, 50 mM Britton-Robinson, 0.1 mM  $\text{CaCl}_2$ . The pH activity is measured at pH 4.0, 4.5, 5.0, 5.5, 6.0, 7.0,  
30 8.0, 9.5, 9.5, 10, and 10.5, using the Phadebas assay described above.

#### Determination Of AGU Activity and As AGU/mg

One Novo Amyloglucosidase Unit (AGU) is defined as the  
35 amount of enzyme, which hydrolyzes 1 micromole maltose per minute at 37°C and pH 4.3. A detailed description of the

analytical method (AEL-SM-0131) is available on request from Novo Nordisk.

The activity is determined as AGU/ml by a method modified after (AEL-SM-0131) using the Glucose GOD-Perid kit from  
5 Boehringer Mannheim, 124036. Standard: AMG-standard, batch 7-1195, 195 AGU/ml.

375 microL substrate (1% maltose in 50 mM Sodium acetate, pH 4.3) is incubated 5 minutes at 37°C. 25 microL enzyme diluted in sodium acetate is added. The reaction is stopped  
10 after 10 minutes by adding 100 microL 0.25 M NaOH. 20 microL is transferred to a 96 well microtitre plate and 200 microL GOD-Perid solution is added. After 30 minutes at room temperature, the absorbance is measured at 650 nm and the activity calculated in AGU/ml from the AMG-standard.

15 The specific activity in AGU/mg is then calculated from the activity (AGU/ml) divided with the protein concentration (mg/ml).

#### EXAMPLES

20

##### EXAMPLE 1

Construction of Termamyl variants in accordance with the invention

Termamyl (*B. licheniformis* alpha-amylase SEQ ID NO: 4) is  
25 expressed in *B. subtilis* from a plasmid denoted pDN1528. This plasmid contains the complete gene encoding Termamyl, *amyL*, the expression of which is directed by its own promoter. Further, the plasmid contains the origin of replication, *ori*, from plasmid pUB110 and the *cat* gene from plasmid pC194 conferring  
30 resistance towards chloramphenicol. pDN1528 is shown in Fig. 9 of WO 96/23874. A specific mutagenesis vector containing a major part of the coding region of SEQ ID NO: 3 was prepared. The important features of this vector, denoted pJeEN1, include an origin of replication derived from the pUC plasmids, the *cat*  
35 gene conferring resistance towards chloramphenicol, and a frameshift-containing version of the *bla* gene, the wild type of which normally confers resistance towards ampicillin ( $\text{amp}^R$ )

phenotype). This mutated version results in an amp<sup>s</sup> phenotype. The plasmid pJeEN1 is shown in Fig. 10 of WO 96/23874, and the *E. coli* origin of replication, *ori*, *bla*, *cat*, the 5'-truncated version of the Termamyl amylase gene, and selected restriction sites are indicated on the plasmid.

Mutations are introduced in *amyL* by the method described by Deng and Nickoloff (1992, Anal. Biochem. 200, pp. 81-88) except that plasmids with the "selection primer" (primer #6616; see below) incorporated are selected based on the amp<sup>R</sup> phenotype of transformed *E. coli* cells harboring a plasmid with a repaired *bla* gene, instead of employing the selection by restriction enzyme digestion outlined by Deng and Nickoloff. Chemicals and enzymes used for the mutagenesis were obtained from the Chameleon<sup>®</sup> mutagenesis kit from Stratagene (catalogue number 200509).

After verification of the DNA sequence in variant plasmids, the truncated gene, containing the desired alteration, is subcloned into pDN1528 as a *Pst*I-*Eco*RI fragment and transformed into the protease- and amylase-depleted *Bacillus subtilis* strain SHA273 (described in WO92/11357 and WO95/10603) in order to express the variant enzyme.

The Termamyl variant V54W was constructed by the use of the following mutagenesis primer (written 5' to 3', left to right):

PG GTC GTA GGC ACC GTA GCC CCA ATC CGC TTG (SEQ ID NO: 9)

The Termamyl variant A52W + V54W was constructed by the use of the following mutagenesis primer (written 5' to 3', left to right):

PG GTC GTA GGC ACC GTA GCC CCA ATC CCA TTG GCT CG (SEQ ID NO: 10)

Primer #6616 (written 5' to 3', left to right; P denotes a 5' phosphate):

P CTG TGA CTG GTG AGT ACT CAA CCA AGT C (SEQ ID NO: 11)

The Termamyl variant V54E was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PGG TCG TAG GCA CCG TAG CCC TCA TCC GCT TG (SEQ ID NO: 12)

The Termamyl variant V54M was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PGG TCG TAG GCA CCG TAG CCC ATA TCC GCT TG (SEQ ID NO: 13)

5 The Termamyl variant V54I was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PGG TCG TAG GCA CCG TAG CCA ATA TCC GCT TG (SEQ ID NO: 14)

10 The Termamyl variants Y290E and Y290K were constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PGC AGC ATG GAA CTG CTY ATG AAG AGG CAC GTC AAA C (SEQ ID NO:15)

15 Y represents an equal mixture of C and T. The presence of a codon encoding either Glutamate or Lysine in position 290 was verified by DNA sequencing.

The Termamyl variant N190F was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

20 PCA TAG TTG CCG AAT TCA TTG GAA ACT TCC C (SEQ ID NO: 16)

The Termamyl variant N188P+N190F was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PCA TAG TTG CCG AAT TCA GGG GAA ACT TCC CAA TC (SEQ ID NO: 17)

25 The Termamyl variant H140K+H142D was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PCC GCG CCC CGG GAA ATC AAA TTT TGT CCA GGC TTT AAT TAG (SEQ ID NO: 18)

30 The Termamyl variant H156Y was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PCA AAA TGG TAC CAA TAC CAC TTA AAA TCG CTG (SEQ ID NO: 19)

35 The Termamyl variant A181T was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PCT TCC CAA TCC CAA GTC TTC CCT TGA AAC (SEQ ID NO: 20)

The Termamyl variant A209V was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PCTT AAT TTC TGC TAC GAC GTC AGG ATG GTC ATA ATC (SEQ ID NO: 21)

The Termamyl variant Q264S was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PCG CCC AAG TCA TTC GAC CAG TAC TCA GCT ACC GTA AAC (SEQ ID NO: 22)

The Termamyl variant S187D was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PGC CGT TTT CAT TGT CGA CTT CCC AAT CCC (SEQ ID NO: 23)

The Termamyl variant DELTA(K370-G371-D372) (i.e., deleted of amino acid residues nos. 370, 371 and 372) was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PGG AAT TTC GCG CTG ACT AGT CCC GTA CAT ATC CCC (SEQ ID NO: 24)

The Termamyl variant DELTA(D372-S373-Q374) was constructed by the use of the following mutagenesis primer (written 5'-3', left to right):

PGG CAG GAA TTT CGC GAC CTT TCG TCC CGT ACA TAT C (SEQ ID NO: 25)

The Termamyl variants A181T and A209V were combined to A181T+A209V by digesting the A181T containing pDN1528-like plasmid (i.e., pDN1528 containing within *amyL* the mutation resulting in the A181T alteration) and the A209V-containing pDN1528-like plasmid (i.e., pDN1528 containing within *amyL* the mutation resulting in the A209V alteration) with restriction enzyme *ClaI* which cuts the pDN1528-like plasmids twice resulting in a fragment of 1116 bp and the vector-part (i.e. contains the plasmid origin of replication) of 3850 bp. The fragment containing the A209V mutation and the vector part containing the A181T mutation were purified by QIAquick gel extraction kit (purchased from QIAGEN) after separation on an



agarose gel. The fragment and the vector were ligated and transformed into the protease and amylase depleted *Bacillus subtilis* strain referred to above. Plasmid from amy<sup>+</sup> (clearing zones on starch containing agar-plates) and chloramphenicol resistant transformants were analysed for the presence of both mutations on the plasmid.

In a similar way as described above, H156Y and A209V were combined utilizing restriction endonucleases Acc65I and EcoRI, giving H156Y+A209V.

H156Y +A209V and A181T+A209V were combined into H156Y+A181T+A209V by the use of restriction endonucleases Acc65I and HindIII.

The 35 N-terminal residues of the mature part of Termamyl variant H156Y+ A181T+A209V were substituted by the 33 N-terminal residues of the *B. amyloliquefaciens* alpha-amylase (SEQ ID NO: 4) (which in the present context is termed BAN) by a SOE-PCR approach (Higuchi et al. 1988, Nucleic Acids Research 16:7351) as follows:

Primer 19364 (sequence 5'-3'): CCT CAT TCT GCA GCA GCA GCC GTA AAT GGC ACG CTG (SEQ ID NO: 26)

Primer 19362: CCA GAC GGC AGT AAT ACC GAT ATC CGA TAA ATG TTC CG (SEQ ID NO: 27)

Primer 19363: CGG ATA TCG GTA TTA CTG CCG TCT GGA TTC (SEQ ID NO: 28)

Primer 1C: CTC GTC CCA ATC GGT TCC GTC (SEQ ID NO: 29)

A standard PCR, polymerase chain reaction, was carried out using the Pwo thermostable polymerase from Boehringer Mannheim according to the manufacturer's instructions and the temperature cyclus: 5 minutes at 94°C, 25 cycles of (94°C for 30 seconds, 50°C for 45 seconds, 72°C for 1 minute), 72°C for 10 minutes.

An approximately 130 bp fragment was amplified in a first PCR denoted PCR1 with primers 19364 and 19362 on a DNA fragment containing the gene encoding the *B. amyloliquefaciens* alpha-amylase.

An approximately 400 bp fragment was amplified in another PCR denoted PCR2 with primers 19363 and 1C on template

pDN1528.

PCR1 and PCR2 were purified from an agarose gel and used as templates in PCR3 with primers 19364 and 1C, which resulted in a fragment of approximately 520 bp. This fragment thus  
5 contains one part of DNA encoding the N-terminus from BAN fused to a part of DNA encoding Termamyl from the 35th amino acid.

The 520 bp fragment was subcloned into a pDN1528-like plasmid (containing the gene encoding Termamyl variant H156Y+  
10 A181T+A209V) by digestion with restriction endonucleases *Pst*I and *Sac*II, ligation and transformation of the *B. subtilis* strain as previously described. The DNA sequence between restriction sites *Pst*I and *Sac*II was verified by DNA sequencing in extracted plasmids from *amy*<sup>+</sup> and chloramphenicol  
15 resistant transformants.

The final construct containing the correct N-terminus from BAN and H156Y+ A181T+A209V was denoted BAN(1-35)+ H156Y+ A181T+A209V.

N190F was combined with BAN(1-35)+ H156Y+ A181T+A209V  
20 giving BAN(1-35)+ H156Y+ A181T+N190F+A209V by carrying out mutagenesis as described above except that the sequence of *amy*L in pJeEN1 was substituted by the DNA sequence encoding Termamyl variant BAN(1-35)+ H156Y+ A181T+A209V

Q264S was combined with BAN(1-35)+ H156Y+ A181T+A209V  
25 giving BAN(1-35)+ H156Y+ A181T+A209V+Q264S by carrying out mutagenesis as described above except that the sequence of *amy*L in pJeEN was substituted by the the DNA sequence encoding Termamyl variant BAN(1-35)+ H156Y+ A181T+A209V

BAN(1-35)+ H156Y+ A181T+A209V+Q264S and BAN(1-35)+ H156Y+  
30 A181T+N190F+A209V were combined into BAN(1-35)+ H156Y+ A181T+N190F+A209V+Q264S utilizing restriction endonucleases *Bsa*HI (*Bsa*HI site was introduced close to the A209V mutation) and *Pst*I.

I201F was combined with BAN(1-35)+ H156Y+  
35 A181T+N190F+A209V+Q264S giving BAN(1-35)+ H156Y+ A181T+N190F+A209V+Q264S+I201F (SEQ ID NO: 2) by carrying out mutagenesis as described above. The mutagenesis primer AM100

was used, introduced the I201F substitution and removed simultaneously a Cla I restriction site, which facilitates easy pin-pointing of mutants.

5 primer AM100:

5'GATGTATGCCGACTTCGATTATGACC 3' (SEQ ID NO: 30)

## EXAMPLE 2

10 Construction of Termamyl-like alpha-amylase variants with an altered cleavage pattern according to the invention

The variant of the thermostable *B. licheniformis* alpha-amylase consisting comprising the 445 C-terminal amino acid residues of the *B. licheniformis* alpha-amylase shown in SEQ ID NO: 4 and the 37 N-terminal amino acid residues of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 15 6, and further comprising the following mutations:

H156Y+A181T+N190F+A209V+Q264S+I201F (the construction of this variant is described in Example 1, and the amino acid sequence shown in SEQ ID NO: 2) has a reduced capability of cleaving an 20 substrate close to the branching point.

In an attempt to further improve the reduced capability of cleaving an substrate close to the branching point of said alpha-amylase variant site directed mutagenesis was carried out using the Mega-primer method as described by Sarkar and 25 Sommer, 1990 (BioTechniques 8: 404-407):

Construction of LE313: BAN/Termamyl hybrid + H156Y+A181T+N190F+ A209V+Q264S+V54N:

30 Gene specific primer 27274 and mutagenic primer AM115 are used to amplify by PCR an approximately 440 bp DNA fragment from a pDN1528-like plasmid (harbouring the BAN(1-35)+H156Y+A181T+N190F+I201F+A209V+Q264S mutations in the gene encoding the amylase from SEQ ID NO: 4).

The 440 bp fragment is purified from an agarose gel and used 35 as a Mega-primer together with primer 113711 in a second PCR carried out on the same template.

The resulting approximately 630 bp fragment is digested

with restriction enzymes EcoR V and Acc65 I and the resulting approximately 370 bp DNA fragment is purified and ligated with the pDN1528-like plasmid digested with the same enzymes. Competent *Bacillus subtilis* SHA273 (amylase and protease low) cells are transformed with the ligation and Chloramphenicol resistant transformants are checked by DNA sequencing to verify the presence of the correct mutations on the plasmid.

Primer 27274:

10 5' CATAGTTGCCGAATTCATTGGAACTTCCC 3' (SEQ ID NO: 31)

Primer 1B:

5' CCGATTGCTGACGCTGTTATTTGC 3' (SEQ ID NO: 32)

15 primer AM115:

5' GCCAAGCGGATAACGGCTACGGTGC 3' (SEQ ID NO:33)

Construction of LE314: BAN/Termamyl hybrid + H156Y+A181T+N190F+ A209V+Q264S + A52S is carried out in a similar way, except that mutagenic primer AM116 is used.

AM116:

5' GAACGAGCCAATCGGACGTGGGCTACGG 3' (SEQ ID NO: 34)

25 Construction of LE315: BAN/Termamyl hybrid + H156Y+A181T+N190F+ A209V+Q264S + A52S+V54N is carried out in a similar way, except that mutagenic primer AM117 is used.

AM117:

30 5' GGAACGAGCCAATCGGATAACGGCTACGGTGC 3' (SEQ ID NO: 35)

Construction of LE316: BAN/Termamyl hybrid + H156Y+A181T+N190F+ A209V+Q264S + T49L is carried out in a similar way, except that mutagenic primer AM118 is used.

35

AM118:

5' GCATATAAGGGACTGAGCCAAGCGG 3' (SEQ ID NO: 36)

Construction LE317: BAN/Termamyl hybrid + H156Y+A181T+N190F+ A209V+Q264S + T49L+G107A is carried out in a similar way, except that mutagenic primer AM118 and mutagenic primer AM119 are used simultaneously.

AM119:

5' CAACCACAAAGCCGGCGCTGATGCG 3' (SEQ ID NO: 37)

Construction of LE318: BAN/Termamyl hybrid + H156Y+A181T+N190F+ A209V+Q264S + A52S+V54N+T49L+G107A is carried out in a similar way, except that mutagenic primer AM120 and mutagenic primer AM119 are used simultaneously.

AM120:

5' GCATATAAGGGACTGAGCCAATCGGATAACGGCTACGGTGC 3' (SEQ ID NO: 38)

Construction of LE 319: BAN/Termamyl hybrid + H156Y+A181T+N190F+ A209V+Q264S + A52S+V54N+T49L is carried out in a similar way, except that mutagenic primer AM120 is used.

Construction of LE320: BAN/Termamyl hybrid + H156Y+A181T+N190F+ A209V+Q264S + G107A is carried out in a similar way, except that mutagenic primer AM119 is used.

Construction of LE322: BAN/Termamyl hybrid + H156Y+A181T+N190F+A209V+Q264S + Q51R+A52S is carried out in a similar way, except that mutagenic primer AM121 is used.

AM121:

5' GAACGAGCCGATCGGACGTGGGCTACGG 3' (SEQ ID NO:39)

Construction of LE323: BAN/Termamyl hybrid + H156Y+A181T+N190F+ A209V+Q264S + A52N is carried out in a similar way, except that mutagenic primer AM122 is used.

AM122:

5' GAACGAGCCAAAACGACGTGGGCTACGG 3' (SEQ ID NO: 40)

35

### EXAMPLE 3

Testing of LE429 variants (saccharification)

The standard reaction conditions were:

Substrate concentration    30 % w/w  
 Temperature                60°C  
 Initial pH (at 60°C)       5.5  
 Enzyme dosage  
     Glucoamylase            0.18 AGU/g DS  
     Pullulanase             0.06 PUN/g DS  
     Alpha-amylase          10 micro g enzyme/g DS

Dextrozyme™ E was used to provide glucoamylase and pullulanase activities

Substrates for saccharification were prepared by  
 5 dissolving common corn starch in deionized water and adjusting  
 the dry substance to approximately 30% w/w. The pH was  
 adjusted to 5.5 (measured at 60°C), and aliquots of substrate  
 corresponding to 10 g dry weight were transferred to blue cap  
 glass flasks.

10 The flasks were then placed in a shaking water bath  
 equilibrated at 60°C, and the enzymes added. The pH was  
 readjusted to 5.5 where necessary. The samples were taken  
 after 48 hours of saccharification; the pH was adjusted to  
 about 3.0, and then heated in a boiling water bath for 15  
 15 minutes to inactivate the enzymes. After cooling, the samples  
 were treated with approximately 0.1 g mixed bed ion exchange  
 resin (BIO-RAD 501 X8 (D)) for 30 minutes on a rotary mixer to  
 remove salts and soluble N. After filtration, the  
 carbohydrate composition was determined by HPLC. The following  
 20 results were obtained:

The parent alpha-amylase for the variants is LE429.

Added Alpha-amylase Variants	DP <sub>1</sub>	DP <sub>2</sub>	DP <sub>3</sub>	SPEC. ACT. (NU/mg)
V54N	96.1	1.75	1.18	8200
A52S	95.9	1.80	1.11	18800
A52S+V54N	96.3	1.84	1.08	10000

T49L	96.3	1.77	1.11	12300
T49L+G107A	96.4	1.87	0.72	13600
A52S+V54N+T49L+G107A	80.5	2.55	0.43	10000
A52S+V54N+T49L	95.8	1.76	0.84	8400
G107A	94.4	1.89	1.04	19600
Q51R+A52S	95.9	1.77	1.27	16500
A52N	95.5	1.89	1.56	17600
LE174 (CONTROL)	95.9/ 95.8	1.87/ 1.83	1.17/ 1.35	16000

Compared with the control, the presence of an active alpha-amylase variant of the invention during liquefaction results in decreased panose levels (DP3).

5 Especially the T49L+G107A variant of LE429 and the A52S+V54N+T49L variant of LE429, respectively, result in a drastically decreased panose level (DP<sub>3</sub>). If these alpha-amylase variants are used for starch liquefaction, it will not be necessary to inactivate the enzyme before the commencement  
10 of saccharification.

#### Example 4

##### Liquefaction and saccharification of LE429 variants

The experiment in Example 3 was repeated for a number of  
15 other LE429 variants under the same conditions.

The result is shown below:

Variant/sugar profile	DP1	DP2	DP3	DP4+
T49V+G107A	95.9%	1.72%	1.27%	1.11%
20 T49Y+G107A	95.3%	1.73%	1.29%	1.65%
T49N+G107A	95.7%	1.64%	1.51%	1.18%
T49L+A52S+G107A	95.7%	1.73%	0.95%	1.67%
T49L+A52T+G107A	95.8%	1.66%	1.03%	1.48%
T49L+A52F+G107A	95.7%	1.69%	1.16%	1.42%
25 T49L+A52L+G107A	95.5%	1.70%	1.40%	1.38%
T49L+A52I+G107A	95.9%	1.72%	1.31%	1.07%
T49L+A52V+G107A	94.7%	1.69%	1.16%	2.44%

T49L+A52V+G107A+A111V	94.5%	1.75%	0.72%	2.99%
LE429	94.9%	1.71%	1.85%	1.51%

**Example 5**

- 5       The experiment in Example 3 was repeated for a number of LE429 variants, except that the liquefaction was carried out at 95°C, pH 6.0 and the saccharification at 60°C, pH 4.5, 40 ppm CaCl<sub>2</sub>, followed by inactivation. The variant referred to below are LE429 variant. The results found are as follows:

10

Variant/sugar profile	DP4+	DP3	DP2	DP1
T49F	1.15	0.92	1.83	96.12
T49D+G107A	0.84	1.03	1.82	96.3
T49I+G107A	0.97	0.64	1.84	96.55
15 T49L+G107A	0.96	0.81	1.82	96.42
T49L+A52S+G107A	1.37	0.75	1.88	96.01
T49L+A52T+G107A	0.87	0.81	1.8	96.52
T49L+A52F+G107A	0.98	0.83	1.87	96.31
T49V+G107A	0.65	0.8	2.13	96.43
20 T49Y+G107A	0.83	0.94	1.89	96.35
LE429	1.16	1.21	1.77	95.87



## REFERENCES CITED

- Klein, C., et al., *Biochemistry* 1992, 31, 8740-8746,  
Mizuno, H., et al., *J. Mol. Biol.* (1993) 234, 1282-1283,  
5 Chang, C., et al., *J. Mol. Biol.* (1993) 229, 235-238,  
Larson, S.B., *J. Mol. Biol.* (1994) 235, 1560-1584,  
Lawson, C.L., *J. Mol. Biol.* (1994) 236, 590-600,  
Qian, M., et al., *J. Mol. Biol.* (1993) 231, 785-799,  
Brady, R.L., et al., *Acta Crystallogr. sect. B*, 47, 527-535,  
10 Swift, H.J., et al., *Acta Crystallogr. sect. B*, 47, 535-544  
A. Kadziola, Ph.D. Thesis: "An alpha-amylase from Barley and its  
Complex with a Substrate Analogue Inhibitor Studied by X-ray  
Crystallography", Department of Chemistry University of  
Copenhagen 1993  
15 MacGregor, E.A., *Food Hydrocolloids*, 1987, Vol.1, No. 5-6, p.  
B. Diderichsen and L. Christiansen, Cloning of a maltogenic  
amylase from *Bacillus stearothermophilus*, *FEMS Microbiol. let-  
ters*: 56: pp. 53-60 (1988)  
Hudson et al., *Practical Immunology*, Third edition (1989),  
20 Blackwell Scientific Publications,  
Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd  
Ed., Cold Spring Harbor, 1989  
S.L. Beaucage and M.H. Caruthers, Tetrahedron Letters 22, 1981,  
pp. 1859-1869  
25 Matthes et al., The EMBO J. 3, 1984, pp. 801-805.  
R.K. Saiki et al., Science 239, 1988, pp. 487-491.  
Morinaga et al., (1984, *Biotechnology* 2:646-639)  
Nelson and Long, Analytical Biochemistry 180, 1989, pp. 147-151  
Hunkapiller et al., 1984, *Nature* 310:105-111  
30 R. Higuchi, B. Krummel, and R.K. Saiki (1988). A general method  
of *in vitro* preparation and specific mutagenesis of DNA frag-  
ments: study of protein and DNA interactions. *Nucl. Acids Res.*  
16:7351-7367.  
Dubnau et al., 1971, *J. Mol. Biol.* 56, pp. 209-221.  
35 Gryczan et al., 1978, *J. Bacteriol.* 134, pp. 318-329.  
S.D. Erlich, 1977, Proc. Natl. Acad. Sci. 74, pp. 1680-1682.  
Boel et al., 1990, Biochemistry 29, pp. 6244-6249.

Sarkar and Sommer, 1990, BioTechniques 8, pp. 404-407.

## CLAIMS

1. A variant of a parent Termamyl-like alpha-amylase, comprising an alteration at one or more positions selected from the group of:

W13, G48, T49, S50, Q51, A52, D53, V54, G57, G107, G108, A111, S168, M197, wherein (a) the alteration(s) are independently

(i) an insertion of an amino acid downstream of the amino acid which occupies the position,

(ii) a deletion of the amino acid which occupies the position, or

(iii) a substitution of the amino acid which occupies the position with a different amino acid,

(b) the variant has alpha-amylase activity and (c) each position corresponds to a position of the amino acid sequence of the parent Termamyl-like alpha-amylase having the amino acid sequence of SEQ ID NO: 4.

2. The variant of claims 1, comprises a mutation in a position corresponding to at least one of the following mutations in the amino acid sequence shown in SEQ ID NO: 4:

V54N, A52S, A52S+V54N, T49L, T49+G107A, A52S+V54N+T49L+G107A, A52S+V54N+T49L, G107A, Q51R, Q51R+A52S, A52N; or

T49F+G107A, T49V+G107A, T49D+G107A, T49Y+G107A, T49S+G107A, T49N+G107A, T49I+G107A, T49L+A52S+G107A, T49L+A52T+G107A, T49L+A52F+G107A, T49L+A52L+G107A, T49L+A52I+G107A, T49L+A52V+G107A; or

T49V, T49I, T49D, T49N, T49S, T49Y, T49F, T49W, T49M, T49E, T49Q, T49K, T49R, A52T, A52L, A52I, A52V, A52M, A52F, A52Y, A52W, V54M, G107V, G07I, G107L, G107C.

3. The variant of claims 1 or 2, comprising a mutation in a position corresponding to at least one of the following mutations in the amino acid sequence shown in SEQ ID NO: 4:

W13F, L, I, V, Y, A;

G48A, V, S, T, I, L;

\*48aD or \*48aY (i.e., insertion of D or Y);

T49X;

\*49aX (i.e., insertion of any amino acid residue)

S50X, in particular D,Y,L,T,V,I;

Q51R,K;

5 A52X, in particular A52S,N,T,F,L,I,V;

D53E,Q,Y,I,N,S,T,V,L;

V54X, in particular V54I,N,W,Y,F,L;

G57S,A,V,L,I,F,Y,T;

G107X, in particular G107A,V,S,T,I,L,C;

10 G108X, in particular G108A,V,S,T,I,L;

A111V,I,L;

S168Y;

M197X, in particular Y,F,L,I,T,A,G.

15

4. The variant of any of claims 1-3, comprises the following mutations corresponding to at least one of the following mutations in the amino acid sequence shown in SEQ ID NO: 4:

T49X+A52X+V54N/I/L/Y/F/W+G107A.

20

5. The variant of claims 1-4, further comprising G108A.

6. The variant of claim 1-5, comprises the following mutations corresponding to at least one of the following mutations in the amino acid sequence shown in SEQ ID NO: 4:

25

T49L+G107A;

T49I+G107A;

T49L+G107A+V54I;

T49I+G107A+V54I;

30

A52S+V54N+T49L+G107A;

A52S+V54I+T49L+G107A;

A52S+T49L+G107A;

A52T+T49L+G107A;

A52S+V54N+T49I+G107A;

35

A52S+V54I+T49I+G107A;

A52S+T49I+G107A;

T49L+G108A;

T49I+G108A;  
T49L+G108A+V54I;  
T49I+G108A+V54I.

- 5 7. A variant of any of claims 1-6, wherein said variant has a reduced capability of cleaving an oligo-saccharide substrate close to the branching point as compared to the parent alpha-amylase.
- 10 8. A variant of any of claims 1-7, which further exhibits improved substrate specificity and/or improved specific activity relative to the parent Termamyl-like alpha-amylase.
- 15 9. A variant of any of claims 1-8, wherein the parent alpha-amylase is a hybrid alpha-amylase of SEQ ID NO: 4 and SEQ ID NO: 6.
- 20 10. The variant of any of claims 1-9, wherein the parent hybrid alpha-amylase is a hybrid alpha-amylase comprising the 445 C-terminal amino acid residues of the *B. licheniformis* alpha-amylase shown in SEQ ID NO: 4 and the 37 N-terminal amino acid residues of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6.
- 25 11. The variant of any of claims 1-10, wherein the parent hybrid Termamyl-like alpha-amylase further has the following mutations: H156Y+A181T+N190F+A209V+Q264S (using the numbering in SEQ ID NO: 4) or LE174.
- 30 12. The variant of any of claims 1-11, wherein the parent hybrid Termamyl-like alpha-amylase further has the following mutations: H156Y+A181T+N190F+A209V+Q264S+I201F (using the numbering of SEQ ID NO: 4) or LE429.
- 35 13. A DNA construct comprising a DNA sequence encoding an alpha-amylase variant according to any of claims 1-12.

14. A recombinant expression vector which carries a DNA construct according to claim 13.

15. A cell which is transformed with a DNA construct according to claim 13 or a vector according to claim 14.

16. A cell of claim 9, which is a microorganism, in particular a bacterium or a fungus, such as a gram positive bacterium such as *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*,  
10 *Bacillus brevis*, *Bacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*, *Bacillus circulans*, *Bacillus lautus* or *Bacillus thuringiensis*.

17. A composition comprising:

15 (i) a mixture of the alpha-amylase from *B. licheniformis* having the sequence shown in SEQ ID NO: 4 with one or more variants of claims 1-12 derived from (as the parent Termamyl-like alpha-amylase) the *B. stearothermophilus* alpha-amylase having the sequence shown in SEQ ID NO: 8; or

20 (ii) a mixture of the alpha-amylase from *B. stearothermophilus* having the sequence shown in SEQ ID NO: 8 with one or more variants of claims 1-12 derived from one or more other parent Termamyl-like alpha-amylases; or

(iii) a mixture of one or more variants of claim 1-12 derived  
25 from (as the parent Termamyl-like alpha-amylase) the *B. stearothermophilus* alpha-amylase having the sequence shown in SEQ ID NO: 8 with one or more variants according to the invention derived from one or more other parent Termamyl-like alpha-amylases.

30

18. A composition comprising:

a mixture of one or more variants of claims 1-12 derived from (as the parent Termamyl-like alpha-amylase) the *B. stearothermophilus* alpha-amylase having the sequence shown in  
35 SEQ ID NO: 8 and a Termamyl-like alpha-amylase derived from the *B. licheniformis* alpha-amylase having the sequence shown in SEQ ID NO: 4.

19. A composition comprising:

a mixture of one or more variants of claims 1-12 derived from  
(as the parent Termamyl-like alpha-amylase) the *B.*  
5 *stearotherophilus* alpha-amylase having the sequence shown in  
SEQ ID NO: 8 and a hybrid alpha-amylase comprising a part of the  
*B. amyloliquefaciens* alpha-amylase shown in SEQ ID NO: 6 and a  
part of the *B. licheniformis* alpha-amylase shown in SEQ ID NO:  
4.

10

20. A composition comprising:

a mixture of one or more variants of claims 1-12 derived from  
(as the parent Termamyl-like alpha-amylase) a hybrid alpha-  
amylase comprising a part of the *B. amyloliquefaciens* alpha-  
15 amylase shown in SEQ ID NO: 6 and a part of the *B. licheniformis*  
alpha-amylase shown in SEQ ID NO: 4.

21. A composition of claim 20, wherein the hybrid alpha-amylase  
is a hybrid alpha-amylase comprising the 445 C-terminal amino  
20 acid residues of the *B. licheniformis* alpha-amylase shown in SEQ  
ID NO: 4 and the 37 N-terminal amino acid residues of the alpha-  
amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6.

22. A composition of claim 21, wherein the hybrid alpha-amylase  
25 further has the following mutations:

H156Y+A181T+N190F+A209V+Q264S (using the numbering in SEQ ID NO:  
4) or LE174.

23. A composition of claim 21, wherein the hybrid alpha-amylase  
30 further has the following mutations:

H156Y+A181T+N190F+A209V+Q264S+I201F as shown in SEQ ID NO: 2 or  
LE429.

24. A method for generating a variant of a parent Termamyl-like  
35 alpha-amylase, which variant exhibits a reduced capability of  
cleaving a substrate close to the branching point, and further  
exhibits improved substrate specificity and/or improved

specific activity relative to the parent, the method comprising:

(a) subjecting a DNA sequence encoding the parent Termamyl-like alpha-amylase to random mutagenesis,

5 (b) expressing the mutated DNA sequence obtained in step (a) in a host cell, and

(c) screening for host cells expressing a mutated alpha-amylase which has increased stability at low pH and low calcium concentration relative to the parent alpha-amylase.

10

25. Use of an alpha-amylase variant of any of claims 1-12 or a composition of any of claims 17-23 for starch liquefaction; in detergent composition, such as laundry, dish washing and hard surface cleaning compositions; ethanol production, such as fuel,  
15 drinking and industrial ethanol production; desizing of textiles, fabrics or garments.



## SEQUENCE LISTING

5 <110> Novo Nordisk A/S  
 <120>  
 <130>  
 10 <160> 40  
 <170> PatentIn Ver. 2.1  
 <210> 1  
 15 <211> 1443  
 <212> DNA  
 <213> Bacillus amyloliquefaciens  
 <220>  
 20 <221> CDS  
 <222> (1)..(1443)  
 <400> 1  
 25 gta aat ggc acg ctg atg cag tat ttt gaa tgg tat acg ccg aac gac 48  
 Val Asn Gly Thr Leu Met Gln Tyr Phe Glu Trp Tyr Thr Pro Asn Asp  
 1 5 10 15  
 ggc cag cat tgg aaa cga ttg cag aat gat ggc gaa cat tta tgc gat 96  
 Gly Gln His Trp Lys Arg Leu Gln Asn Asp Ala Glu His Leu Ser Asp  
 30 20 25 30  
 atc ggt att act gcc gtc tgg att ccc ccg gca tat aag gga acg agc 144  
 Ile Gly Ile Thr Ala Val Trp Ile Pro Pro Ala Tyr Lys Gly Thr Ser  
 35 35 40 45  
 caa gcg gat gtg ggc tac ggt gct tac gac ctt tat gat tta ggg gag 192  
 Gln Ala Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr Asp Leu Gly Glu  
 50 55 60  
 40 ttt cat caa aaa ggg acg gtt cgg aca aag tac ggc aca aaa gga gag 240  
 Phe His Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly Thr Lys Gly Glu  
 65 70 75 80  
 ctg caa tct gcg atc aaa agt ctt cat tcc cgc gac att aac gtt tac 288  
 Leu Gln Ser Ala Ile Lys Ser Leu His Ser Arg Asp Ile Asn Val Tyr  
 85 90 95  
 ggg gat gtg gtc atc aac cac aaa ggc ggc gct gat gcg acc gaa gat 336  
 Gly Asp Val Val Ile Asn His Lys Gly Gly Ala Asp Ala Thr Glu Asp  
 50 100 105 110  
 gta acc gcg gtt gaa gtc gat ccc gct gac cgc aac cgc gta att tca 384  
 Val Thr Ala Val Glu Val Asp Pro Ala Asp Arg Asn Arg Val Ile Ser  
 115 120 125  
 55 gga gaa cac cta att aaa gcc tgg aca cat ttt cat ttt ccg ggg cgc 432  
 Gly Glu His Leu Ile Lys Ala Trp Thr His Phe His Phe Pro Gly Arg  
 130 135 140  
 60 ggc agc aca tac agc gat ttt aag tgg tat tgg tac cat ttt gac gga 480  
 Gly Ser Thr Tyr Ser Asp Phe Lys Trp Tyr Trp Tyr His Phe Asp Gly  
 145 150 155 160  
 65 acc gat tgg gac gag tcc cga aag ctg aac cgc atc tat aag ttt caa 528  
 Thr Asp Trp Asp Glu Ser Arg Lys Leu Asn Arg Ile Tyr Lys Phe Gln  
 165 170 175

	ggg aag act tgg gat tgg gaa gtt tcc aat gaa ttc ggc aac tat gat	576
	Gly Lys Thr Trp Asp Trp Glu Val Ser Asn Glu Phe Gly Asn Tyr Asp	
	180 185 190	
5	tat ttg atg tat gcc gac ttt gat tat gac cat cct gat gtc gta gca	624
	Tyr Leu Met Tyr Ala Asp Phe Asp Tyr Asp His Pro Asp Val Val Ala	
	195 200 205	
10	gag att aag aga tgg ggc act tgg tat gcc aat gaa ctg caa ttg gac	672
	Glu Ile Lys Arg Trp Gly Thr Trp Tyr Ala Asn Glu Leu Gln Leu Asp	
	210 215 220	
15	ggt ttc cgt ctt gat gct gtc aaa cac att aaa ttt tct ttt ttg cgg	720
	Gly Phe Arg Leu Asp Val Lys His Ile Lys Phe Ser Phe Leu Arg	
	225 230 235 240	
20	gat tgg gtt aat cat gtc agg gaa aaa acg ggg aag gaa atg ttt acg	768
	Asp Trp Val Asn His Val Arg Glu Lys Thr Gly Lys Glu Met Phe Thr	
	245 250 255	
25	gta gct gag tac tgg tcg aat gac ttg ggc gcg ctg gaa aac tat ttg	816
	Val Ala Glu Tyr Trp Ser Asn Asp Leu Gly Ala Leu Glu Asn Tyr Leu	
	260 265 270	
30	aac aaa aca aat ttt aat cat tca gtg ttt gac gtg ccg ctt cat tat	864
	Asn Lys Thr Asn Phe Asn His Ser Val Phe Asp Val Pro Leu His Tyr	
	275 280 285	
35	cag ttc cat gct gca tcg aca cag gga ggc ggc tat gat atg agg aaa	912
	Gln Phe His Ala Ala Ser Gln Gly Gly Tyr Asp Met Arg Lys	
	290 295 300	
40	ttg ctg aac ggt acg gtc gtt tcc aag cat ccg ttg aaa tcg gtt aca	960
	Leu Leu Asn Gly Thr Val Ser Lys His Pro Leu Lys Ser Val Thr	
	305 310 315 320	
45	ttt gtc gat aac cat gat aca cag ccg ggg caa tcg ctt gag tcg act	1008
	Phe Val Asp Asn His Asp Thr Gln Pro Gly Gln Ser Leu Glu Ser Thr	
	325 330 335	
50	gtc caa aca tgg ttt aag ccg ctt gct tac gct ttt att ctc aca agg	1056
	Val Gln Thr Trp Phe Lys Pro Leu Ala Tyr Ala Phe Ile Leu Thr Arg	
	340 345 350	
55	gaa tct gga tac cct cag gtt ttc tac ggg gat atg tac ggg acg aaa	1104
	Glu Ser Gly Tyr Pro Gln Val Phe Tyr Gly Asp Met Tyr Gly Thr Lys	
	355 360 365	
60	gga gac tcc cag cgc gaa att cct gcc ttg aaa cac aaa att gaa ccg	1152
	Gly Asp Ser Gln Arg Glu Ile Pro Ala Leu Lys His Lys Ile Glu Pro	
	370 375 380	
65	atc tta aaa gcg aga aaa cag tat gcg tac gga gca cag cat gat tat	1200
	Ile Leu Lys Ala Arg Lys Gln Tyr Ala Tyr Gly Ala Gln His Asp Tyr	
	385 390 395 400	
70	ttc gac cac cat gac att gtc ggc tgg aca agg gaa ggc gac agc tcg	1248
	Phe Asp His His Asp Ile Val Gly Trp Arg Glu Gly Asp Ser Ser	
	405 410 415	
75	gtt gca aat tca ggt ttg gcg gca tta ata aca gac gga ccc ggt ggg	1296
	Val Ala Asn Ser Gly Leu Ala Ala Leu Ile Thr Asp Gly Pro Gly Gly	
	420 425 430	
80	gca aag cga atg tat gtc ggc cgg caa aac gcc ggt gag aca tgg cat	1344
	Ala Lys Arg Met Tyr Val Gly Arg Gln Asn Ala Gly Glu Thr Trp His	
	435 440 445	

gac att acc gga aac cgt tcg gag ccg gtt gtc atc aat tcg gaa ggc 1392  
 Asp Ile Thr Gly Asn Arg Ser Glu Pro Val Val Ile Asn Ser Glu Gly  
 450 455 460  
 5 tgg gga gag ttt cac gta aac ggc ggg tcg gtt tca att tat gtt caa 1440  
 Trp Gly Glu Phe His Val Asn Gly Gly Ser Val Ser Ile Tyr Val Gln  
 465 470 475 480  
 10 aga 1443  
 Arg  
 <210> 2  
 15 <211> 481  
 <212> PRT  
 <213> Bacillus amyloliquefaciens  
 <400> 2  
 20 Val Asn Gly Thr Leu Met Gln Tyr Phe Glu Trp Tyr Thr Pro Asn Asp  
 1 5 10 15  
 Gly Gln His Trp Lys Arg Leu Gln Asn Asp Ala Glu His Leu Ser Asp  
 20 25 30  
 25 Ile Gly Ile Thr Ala Val Trp Ile Pro Pro Ala Tyr Lys Gly Thr Ser  
 35 40 45  
 30 Gln Ala Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr Asp Leu Gly Glu  
 50 55 60  
 Phe His Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly Thr Lys Gly Glu  
 65 70 75 80  
 35 Leu Gln Ser Ala Ile Lys Ser Leu His Ser Arg Asp Ile Asn Val Tyr  
 85 90 95  
 Gly Asp Val Val Ile Asn His Lys Gly Gly Ala Asp Ala Thr Glu Asp  
 100 105 110  
 40 Val Thr Ala Val Glu Val Asp Pro Ala Asp Arg Asn Arg Val Ile Ser  
 115 120 125  
 Gly Glu His Leu Ile Lys Ala Trp Thr His Phe His Phe Pro Gly Arg  
 130 135 140  
 45 Gly Ser Thr Tyr Ser Asp Phe Lys Trp Tyr Trp Tyr His Phe Asp Gly  
 145 150 155 160  
 50 Thr Asp Trp Asp Glu Ser Arg Lys Leu Asn Arg Ile Tyr Lys Phe Gln  
 165 170 175  
 Gly Lys Thr Trp Asp Trp Glu Val Ser Asn Glu Phe Gly Asn Tyr Asp  
 180 185 190  
 55 Tyr Leu Met Tyr Ala Asp Phe Asp Tyr Asp His Pro Asp Val Val Ala  
 195 200 205  
 60 Glu Ile Lys Arg Trp Gly Thr Trp Tyr Ala Asn Glu Leu Gln Leu Asp  
 210 215 220  
 Gly Phe Arg Leu Asp Ala Val Lys His Ile Lys Phe Ser Phe Leu Arg  
 225 230 235 240  
 65 Asp Trp Val Asn His Val Arg Glu Lys Thr Gly Lys Glu Met Phe Thr  
 245 250 255

Val Ala Glu Tyr Trp Ser Asn Asp Leu Gly Ala Leu Glu Asn Tyr Leu  
 260 265 270

5 Asn Lys Thr Asn Phe Asn His Ser Val Phe Asp Val Pro Leu His Tyr  
 275 280 285

Gln Phe His Ala Ala Ser Thr Gln Gly Gly Tyr Asp Met Arg Lys  
 290 295 300

10 Leu Leu Asn Gly Thr Val Val Ser Lys His Pro Leu Lys Ser Val Thr  
 305 310 315 320

Phe Val Asp Asn His Asp Thr Gln Pro Gly Gln Ser Leu Glu Ser Thr  
 325 330 335

15 Val Gln Thr Trp Phe Lys Pro Leu Ala Tyr Ala Phe Ile Leu Thr Arg  
 340 345 350

Glu Ser Gly Tyr Pro Gln Val Phe Tyr Gly Asp Met Tyr Gly Thr Lys  
 355 360 365

Gly Asp Ser Gln Arg Glu Ile Pro Ala Leu Lys His Lys Ile Glu Pro  
 370 375 380

25 Ile Leu Lys Ala Arg Lys Gln Tyr Ala Tyr Gly Ala Gln His Asp Tyr  
 385 390 395 400

Phe Asp His His Asp Ile Val Gly Trp Thr Arg Glu Gly Asp Ser Ser  
 405 410 415

30 Val Ala Asn Ser Gly Leu Ala Ala Leu Ile Thr Asp Gly Pro Gly Gly  
 420 425 430

Ala Lys Arg Met Tyr Val Gly Arg Gln Asn Ala Gly Glu Thr Trp His  
 435 440 445

Asp Ile Thr Gly Asn Arg Ser Glu Pro Val Val Ile Asn Ser Glu Gly  
 450 455 460

40 Trp Gly Glu Phe His Val Asn Gly Gly Ser Val Ser Ile Tyr Val Gln  
 465 470 475 480

Arg

45

&lt;210&gt; 3

&lt;211&gt; 1920

50 &lt;212&gt; DNA

<213> *Bacillus licheniformis*

&lt;220&gt;

&lt;221&gt; CDS

55 &lt;222&gt; (421)..(1872)

&lt;400&gt; 3

cggaagattg gaagtacaaa aataagcaaa agattgtcaa tcatgtcatg agccatgcgg 60

60 gagacggaaa aatcgtctta atgcacgata tttatgcaac gttcgcagat gctgctgaag 120

agattattaa aaagctgaaa gcaaaaggct atcaattggt aactgtatct cagcttgaag 180

aagtgaagaa gcagagaggc tattgaataa atgagtagaa gcgccatatt gccgcttttc 240

65 ttttggaaga aaatataggg aaaatggtac ttgttaaaaa ttcggaatat ttatacaaca 300

```

tcatatgttt cacattgaaa ggggaggaga atcatgaaac aacaaaaacg gctttacgcc 360
cgattgctga cgctgttatt tgcgctcatt ttcttgcctgc ctcattctgc agcagcggcg 420

5  gca aat ctt aat ggg acg ctg atg cag tat ttt gaa tgg tac atg ccc 468
   Ala Asn Leu Asn Gly Thr Leu Met Gln Tyr Phe Glu Trp Tyr Met Pro
   1      5      10      15

10  aat gac ggc caa cat tgg agg cgt ttg caa aac gac tcg gca tat ttg 516
   Asn Asp Gly Gln His Trp Arg Arg Leu Gln Asn Asp Ser Ala Tyr Leu
           20      25      30

15  gct gaa cac ggt att act gcc gtc tgg att ccc ccg gca tat aag gga 564
   Ala Glu His Gly Ile Thr Ala Val Trp Ile Pro Pro Ala Tyr Lys Gly
           35      40      45

20  acg agc caa gcg gat gtg ggc tac ggt gct tac gac ctt tat gat tta 612
   Thr Ser Gln Ala Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr Asp Leu
           50      55      60

25  ggg gag ttt cat caa aaa ggg acg gtt cgg aca aag tac ggc aca aaa 660
   Gly Glu Phe His Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly Thr Lys
   65      70      75      80

30  gga gag ctg caa tct gcg atc aaa agt ctt cat tcc cgc gac att aac 708
   Gly Glu Leu Gln Ser Ala Ile Lys Ser Leu His Ser Arg Asp Ile Asn
           85      90      95

35  gtt tac ggg gat gtg gtc atc aac cac aaa ggc ggc gct gat gcg acc 756
   Val Tyr Gly Asp Val Val Ile Asn His Lys Gly Gly Ala Asp Ala Thr
           100      105      110

40  gaa gat gta acc gcg gtt gaa gtc gat ccc gct gac cgc aac cgc gta 804
   Glu Asp Val Thr Ala Val Glu Val Asp Pro Ala Asp Arg Asn Arg Val
           115      120      125

45  att tca gga gaa cac cta att aaa gcc tgg aca cat ttt cat ttt ccg 852
   Ile Ser Gly Glu His Leu Ile Lys Ala Trp Thr His Phe His Phe Pro
           130      135      140

50  ggg cgc ggc agc aca tac agc gat ttt aaa tgg cat tgg tac cat ttt 900
   Gly Arg Gly Ser Thr Tyr Ser Asp Phe Lys Trp His Trp Tyr His Phe
   145      150      155      160

55  gac gga acc gat tgg gac gag tcc cga aag ctg aac cgc atc tat aag 948
   Asp Gly Thr Asp Trp Asp Glu Ser Arg Lys Leu Asn Arg Ile Tyr Lys
           165      170      175

60  ttt caa gga aag gct tgg gat tgg gaa gtt tcc aat gaa aac ggc aac 996
   Phe Gln Gly Lys Ala Trp Asp Trp Glu Val Ser Asn Glu Asn Gly Asn
           180      185      190

65  tat gat tat ttg atg tat gcc gac atc gat tat gac cat cct gat gtc 1044
   Tyr Asp Tyr Leu Met Tyr Ala Asp Ile Asp Tyr Asp His Pro Asp Val
           195      200      205

70  gca gca gaa att aag aga tgg ggc act tgg tat gcc aat gaa ctg caa 1092
   Ala Ala Glu Ile Lys Arg Trp Gly Thr Trp Tyr Ala Asn Glu Leu Gln
   210      215      220

75  ttg gac ggt ttc cgt ctt gat gct gtc aaa cac att aaa ttt tct ttt 1140
   Leu Asp Gly Phe Arg Leu Asp Ala Val Lys His Ile Lys Phe Ser Phe
   225      230      235      240

80  ttg cgg gat tgg gtt aat cat gtc agg gaa aaa acg ggg aag gaa atg 1188
   Leu Arg Asp Trp Val Asn His Val Arg Glu Lys Thr Gly Lys Glu Met
           245      250      255

```

	ttt acg gta gct gaa tat tgg cag aat gac ttg ggc gcg ctg gaa aac	1236
	Phe Thr Val Ala Glu Tyr Trp Gln Asn Asp Leu Gly Ala Leu Glu Asn	
	260 265 270	
5	tat ttg aac aaa aca aat ttt aat cat tca gtg ttt gac gtg ccg ctt	1284
	Tyr Leu Asn Lys Thr Asn Phe Asn His Ser Val Phe Asp Val Pro Leu	
	275 280 285	
10	cat tat cag ttc cat gct gca tcg aca cag gga ggc ggc tat gat atg	1332
	His Tyr Gln Phe His Ala Ala Ser Thr Gln Gly Gly Gly Tyr Asp Met	
	290 295 300	
15	agg aaa ttg ctg aac ggt acg gtc gtt tcc aag cat ccg ttg aaa tcg	1380
	Arg Lys Leu Leu Asn Gly Thr Val Val Ser Lys His Pro Leu Lys Ser	
	305 310 315 320	
20	ggt aca ttt gtc gat aac cat gat aca cag ccg ggg caa tcg ctt gag	1428
	Val Thr Phe Val Asp Asn His Asp Thr Gln Pro Gly Gln Ser Leu Glu	
	325 330 335	
25	tcg act gtc caa aca tgg ttt aag ccg ctt gct tac gct ttt att ctc	1476
	Ser Thr Val Gln Thr Trp Phe Lys Pro Leu Ala Tyr Ala Phe Ile Leu	
	340 345 350	
30	aca agg gaa tct gga tac cct cag gtt ttc tac ggg gat atg tac ggg	1524
	Thr Arg Glu Ser Gly Tyr Pro Gln Val Phe Tyr Gly Asp Met Tyr Gly	
	355 360 365	
35	acg aaa gga gac tcc cag cgc gaa att cct gcc ttg aaa cac aaa att	1572
	Thr Lys Gly Asp Ser Gln Arg Glu Ile Pro Ala Leu Lys His Lys Ile	
	370 375 380	
40	gaa ccg atc tta aaa gcg aga aaa cag tat gcg tac gga gca cag cat	1620
	Glu Pro Ile Leu Lys Ala Arg Lys Gln Tyr Ala Tyr Gly Ala Gln His	
	385 390 395 400	
45	gat tat ttc gac cac cat gac att gtc ggc ttg aca agg gaa ggc gac	1668
	Asp Tyr Phe Asp His His Asp Ile Val Gly Trp Thr Arg Glu Gly Asp	
	405 410 415	
50	agc tcg gtt gca aat tca ggt ttg gcg gca tta ata aca gac gga ccc	1716
	Ser Ser Val Ala Asn Ser Gly Leu Ala Ala Leu Ile Thr Asp Gly Pro	
	420 425 430	
55	ggg ggc gca aag cga atg tat gtc ggc cgg caa aac gcc ggt gag aca	1764
	Gly Gly Ala Lys Arg Met Tyr Val Gly Arg Gln Asn Ala Gly Glu Thr	
	435 440 445	
60	ttg cat gac att acc gga aac cgt tcg gag ccg gtt gtc atc aat tcg	1812
	Trp His Asp Ile Thr Gly Asn Arg Ser Glu Pro Val Val Ile Asn Ser	
	450 455 460	
65	gaa ggc ttg gga gag ttt cac gta aac ggc ggg tcg gtt tca att tat	1860
	Glu Gly Trp Gly Glu Phe His Val Asn Gly Gly Ser Val Ser Ile Tyr	
	465 470 475 480	
60	ggt caa aga tag aagagcagag aggacggatt tctgaagga aatccgtttt	1912
	Val Gln Arg	
60	tttatattt	1920
65	<210> 4	
	<211> 483	
	<212> PRT	
	<213> Bacillus licheniformis	

<400> 4  
 Ala Asn Leu Asn Gly Thr Leu Met Gln Tyr Phe Glu Trp Tyr Met Pro  
 1 5 10 15  
 5 Asn Asp Gly Gln His Trp Arg Arg Leu Gln Asn Asp Ser Ala Tyr Leu  
 20 25 30  
 10 Ala Glu His Gly Ile Thr Ala Val Trp Ile Pro Pro Ala Tyr Lys Gly  
 35 40 45  
 Thr Ser Gln Ala Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr Asp Leu  
 50 55 60  
 15 Gly Glu Phe His Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly Thr Lys  
 65 70 75 80  
 Gly Glu Leu Gln Ser Ala Ile Lys Ser Leu His Ser Arg Asp Ile Asn  
 85 90 95  
 20 Val Tyr Gly Asp Val Val Ile Asn His Lys Gly Gly Ala Asp Ala Thr  
 100 105 110  
 25 Glu Asp Val Thr Ala Val Glu Val Asp Pro Ala Asp Arg Asn Arg Val  
 115 120 125  
 Ile Ser Gly Glu His Leu Ile Lys Ala Trp Thr His Phe His Phe Pro  
 130 135 140  
 30 Gly Arg Gly Ser Thr Tyr Ser Asp Phe Lys Trp His Trp Tyr His Phe  
 145 150 155 160  
 Asp Gly Thr Asp Trp Asp Glu Ser Arg Lys Leu Asn Arg Ile Tyr Lys  
 165 170 175  
 35 Phe Gln Gly Lys Ala Trp Asp Trp Glu Val Ser Asn Glu Asn Gly Asn  
 180 185 190  
 Tyr Asp Tyr Leu Met Tyr Ala Asp Ile Asp Tyr Asp His Pro Asp Val  
 195 200 205  
 40 Ala Ala Glu Ile Lys Arg Trp Gly Thr Trp Tyr Ala Asn Glu Leu Gln  
 210 215 220  
 45 Leu Asp Gly Phe Arg Leu Asp Ala Val Lys His Ile Lys Phe Ser Phe  
 225 230 235 240  
 Leu Arg Asp Trp Val Asn His Val Arg Glu Lys Thr Gly Lys Glu Met  
 245 250 255  
 50 Phe Thr Val Ala Glu Tyr Trp Gln Asn Asp Leu Gly Ala Leu Glu Asn  
 260 265 270  
 Tyr Leu Asn Lys Thr Asn Phe Asn His Ser Val Phe Asp Val Pro Leu  
 275 280 285  
 55 His Tyr Gln Phe His Ala Ala Ser Thr Gln Gly Gly Gly Tyr Asp Met  
 290 295 300  
 60 Arg Lys Leu Leu Asn Gly Thr Val Val Ser Lys His Pro Leu Lys Ser  
 305 310 315 320  
 Val Thr Phe Val Asp Asn His Asp Thr Gln Pro Gly Gln Ser Leu Glu  
 325 330 335  
 65 Ser Thr Val Gln Thr Trp Phe Lys Pro Leu Ala Tyr Ala Phe Ile Leu  
 340 345 350

```

    Thr Arg Glu Ser Gly Tyr Pro Gln Val Phe Tyr Gly Asp Met Tyr Gly
           355                      360                      365
5   Thr Lys Gly Asp Ser Gln Arg Glu Ile Pro Ala Leu Lys His Lys Ile
    370                      375                      380

    Glu Pro Ile Leu Lys Ala Arg Lys Gln Tyr Ala Tyr Gly Ala Gln His
    385                      390                      395                      400
10  Asp Tyr Phe Asp His His Asp Ile Val Gly Trp Thr Arg Glu Gly Asp
           405                      410                      415

    Ser Ser Val Ala Asn Ser Gly Leu Ala Ala Leu Ile Thr Asp Gly Pro
           420                      425                      430
15  Gly Gly Ala Lys Arg Met Tyr Val Gly Arg Gln Asn Ala Gly Glu Thr
    435                      440                      445

    Trp His Asp Ile Thr Gly Asn Arg Ser Glu Pro Val Val Ile Asn Ser
    450                      455                      460
20  Glu Gly Trp Gly Glu Phe His Val Asn Gly Gly Ser Val Ser Ile Tyr
    465                      470                      475                      480
25  Val Gln Arg

30  <210> 5
    <211> 2604
    <212> DNA
    <213> Bacillus amyloliquefaciens

35  <220>
    <221> -10_signal
    <222> (707)..(712)

    <220>
40  <221> -35_signal
    <222> (729)..(734)

    <220>
    <221> RBS
45  <222> (759)..(762)

    <220>
    <221> sig_peptide
    <222> (770)..(862)
50

    <220>
    <221> mat_peptide
    <222> (863)..(2314)

55  <220>
    <221> terminator
    <222> (2321)..(2376)

    <220>
60  <221> CDS
    <222> (863)..(2314)

    <400> 5
    aagcttcaag cgggtcaatcg gaatgtgcat ctgcgttcat acttaggttt tcacccgcat 60
65  attaagcagg cgtttttgaa ccgtgtgaca gaagctgttc gaaaccccg cgggcgggttt 120

```





	cgc atc ttt aag ttt cgt ggg gaa gga aaa gcg tgg gat tgg gaa gta	1420
	Arg Ile Phe Lys Phe Arg Gly Glu Gly Lys Ala Trp Asp Trp Glu Val	
	175 180 185	
5	tca agt gaa aac ggc aac tat gac tat tta atg tat gct gat gtt gac	1468
	Ser Ser Glu Asn Gly Asn Tyr Asp Tyr Leu Met Tyr Ala Asp Val Asp	
	190 195 200	
10	tac gac cac cct gat gtc gtg gca gag aca aaa aaa tgg ggt atc tgg	1516
	Tyr Asp His Pro Asp Val Val Ala Glu Thr Lys Lys Trp Gly Ile Trp	
	205 210 215	
15	tat gcg aat gaa ctg tca tta gac ggc ttc cgt att gat gcc gcc aaa	1564
	Tyr Ala Asn Glu Leu Ser Leu Asp Gly Phe Arg Ile Asp Ala Ala Lys	
	220 225 230	
20	cat att aaa ttt tca ttt ctg cgt gat tgg gtt cag gcg gtc aga cag	1612
	His Ile Lys Phe Ser Phe Leu Arg Asp Trp Val Gln Ala Val Arg Gln	
	235 240 245 250	
25	gcg acg gga aaa gaa atg ttt acg gtt gcg gag tat tgg cag aat aat	1660
	Ala Thr Gly Lys Glu Met Phe Thr Val Ala Glu Tyr Trp Gln Asn Asn	
	255 260 265	
30	gcc ggg aaa ctg gaa aac tac ttg aat aaa aca agc ttt aat caa tcc	1708
	Ala Gly Lys Leu Glu Asn Tyr Leu Asn Lys Thr Ser Phe Asn Gln Ser	
	270 275 280	
35	gtg ttt gat gtt ccg ctt cat ttc aat tta cag gcg gct tcc tca caa	1756
	Val Phe Asp Val Pro Leu His Phe Asn Leu Gln Ala Ala Ser Ser Gln	
	285 290 295	
40	gga ggc gga tat gat atg agg cgt ttg ctg gac ggt acc gtt gtg tcc	1804
	Gly Gly Gly Tyr Asp Met Arg Arg Leu Leu Asp Gly Thr Val Val Ser	
	300 305 310	
45	agg cat ccg gaa aag gcg gtt aca ttt gtt gaa aat cat gac aca cag	1852
	Arg His Pro Glu Lys Ala Val Thr Phe Val Glu Asn His Asp Thr Gln	
	315 320 325 330	
50	ccg gga cag tca ttg gaa tcg aca gtc caa act tgg ttt aaa ccg ctt	1900
	Pro Gly Gln Ser Leu Glu Ser Thr Val Gln Thr Trp Phe Lys Pro Leu	
	335 340 345	
55	gca tac gcc ttt att ttg aca aga gaa tcc ggt tat cct cag gtg ttc	1948
	Ala Tyr Ala Phe Ile Leu Thr Arg Glu Ser Gly Tyr Pro Gln Val Phe	
	350 355 360	
60	tat ggg gat atg tac ggg aca aaa ggg aca tcg cca aag gaa att ccc	1996
	Tyr Gly Asp Met Tyr Gly Thr Lys Gly Thr Ser Pro Lys Glu Ile Pro	
	365 370 375	
65	tca ctg aaa gat aat ata gag ccg att tta aaa gcg cgt aag gag tac	2044
	Ser Leu Lys Asp Asn Ile Glu Pro Ile Leu Lys Ala Arg Lys Glu Tyr	
	380 385 390	
70	gca tac ggg ccc cag cac gat tat att gac cac ccg gat gtg atc gga	2092
	Ala Tyr Gly Pro Gln His Asp Tyr Ile Asp His Pro Asp Val Ile Gly	
	395 400 405 410	
75	tgg acg agg gaa ggt gac agc tcc gcc gcc aaa tca ggt ttg gcc gct	2140
	Trp Thr Arg Glu Gly Asp Ser Ser Ala Ala Lys Ser Gly Leu Ala Ala	
	415 420 425	
80	tta atc acg gac gga ccc gcc gga tca aag cgg atg tat gcc ggc ctg	2188
	Leu Ile Thr Asp Gly Pro Gly Gly Ser Lys Arg Met Tyr Ala Gly Leu	

430 435 440  
 5 aaa aat gcc ggc gag aca tgg tat gac ata acg ggc aac cgt tca gat 2236  
 Lys Asn Ala Gly Glu Thr Trp Tyr Asp Ile Thr Gly Asn Arg Ser Asp  
 445 450 455  
 10 act gta aaa atc gga tct gac ggc tgg gga gag ttt cat gta aac gat 2284  
 Thr Val Lys Ile Gly Ser Asp Gly Trp Gly Glu Phe His Val Asn Asp  
 460 465 470  
 15 ggg tcc gtc tcc att tat gtt cag aaa taa ggtaataaaa aaacacctcc 2334  
 Gly Ser Val Ser Ile Tyr Val Gln Lys  
 475 480  
 20 aagctgagtg cgggtatcag cttggaggtg cggtttatttt ttcagccgta tgacaaggtc 2394  
 ggcacatcaggt gtgacaaata cggatatgctg gctgtcatag gtgacaaatc cggggttttgc 2454  
 gccgttttggc tttttcacat gtctgatttt tgtataatca acaggcacgg agccggaatc 2514  
 25 tttcgcccttg gaaaaataag cggcgatcgt agctgcttcc aatatggatt gttcatcggg 2574  
 atcgctgctt ttaatcaciaa cgtgggatcc 2604  
 30 <210> 6  
 <211> 483  
 <212> PRT  
 <213> Bacillus amyloliquefaciens  
 35 <400> 6  
 Val Asn Gly Thr Leu Met Gln Tyr Phe Glu Trp Tyr Thr Pro Asn Asp  
 1 5 10 15  
 40 Gly Gln His Trp Lys Arg Leu Gln Asn Asp Ala Glu His Leu Ser Asp  
 20 25 30  
 45 Ile Gly Ile Thr Ala Val Trp Ile Pro Pro Ala Tyr Lys Gly Leu Ser  
 35 40 45  
 50 Gln Ser Asp Asn Gly Tyr Gly Pro Tyr Asp Leu Tyr Asp Leu Gly Glu  
 50 55 60  
 55 Phe Gln Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly Thr Lys Ser Glu  
 65 70 75 80  
 60 Leu Gln Asp Ala Ile Gly Ser Leu His Ser Arg Asn Val Gln Val Tyr  
 85 90 95  
 65 Gly Asp Val Val Leu Asn His Lys Ala Gly Ala Asp Ala Thr Glu Asp  
 100 105 110  
 70 Val Thr Ala Val Glu Val Asn Pro Ala Asn Arg Asn Gln Glu Thr Ser  
 115 120 125  
 75 Glu Glu Tyr Gln Ile Lys Ala Trp Thr Asp Phe Arg Phe Pro Gly Arg  
 130 135 140  
 80 Gly Asn Thr Tyr Ser Asp Phe Lys Trp His Trp Tyr His Phe Asp Gly  
 145 150 155 160  
 85 Ala Asp Trp Asp Glu Ser Arg Lys Ile Ser Arg Ile Phe Lys Phe Arg  
 165 170 175  
 90 Gly Glu Gly Lys Ala Trp Asp Trp Glu Val Ser Ser Glu Asn Gly Asn  
 180 185 190

Tyr Asp Tyr Leu Met Tyr Ala Asp Val Asp Tyr Asp His Pro Asp Val  
 195 200 205  
 5 Val Ala Glu Thr Lys Lys Trp Gly Ile Trp Tyr Ala Asn Glu Leu Ser  
 210 215 220  
 Leu Asp Gly Phe Arg Ile Asp Ala Ala Lys His Ile Lys Phe Ser Phe  
 225 230 235 240  
 10 Leu Arg Asp Trp Val Gln Ala Val Arg Gln Ala Thr Gly Lys Glu Met  
 245 250 255  
 Phe Thr Val Ala Glu Tyr Trp Gln Asn Asn Ala Gly Lys Leu Glu Asn  
 260 265 270  
 15 Tyr Leu Asn Lys Thr Ser Phe Asn Gln Ser Val Phe Asp Val Pro Leu  
 275 280 285  
 20 His Phe Asn Leu Gln Ala Ala Ser Ser Gln Gly Gly Tyr Asp Met  
 290 295 300  
 Arg Arg Leu Leu Asp Gly Thr Val Val Ser Arg His Pro Glu Lys Ala  
 305 310 315 320  
 25 Val Thr Phe Val Glu Asn His Asp Thr Gln Pro Gly Gln Ser Leu Glu  
 325 330 335  
 Ser Thr Val Gln Thr Trp Phe Lys Pro Leu Ala Tyr Ala Phe Ile Leu  
 340 345 350  
 30 Thr Arg Glu Ser Gly Tyr Pro Gln Val Phe Tyr Gly Asp Met Tyr Gly  
 355 360 365  
 35 Thr Lys Gly Thr Ser Pro Lys Glu Ile Pro Ser Leu Lys Asp Asn Ile  
 370 375 380  
 Glu Pro Ile Leu Lys Ala Arg Lys Glu Tyr Ala Tyr Gly Pro Gln His  
 385 390 395 400  
 40 Asp Tyr Ile Asp His Pro Asp Val Ile Gly Trp Thr Arg Glu Gly Asp  
 405 410 415  
 Ser Ser Ala Ala Lys Ser Gly Leu Ala Ala Leu Ile Thr Asp Gly Pro  
 420 425 430  
 45 Gly Gly Ser Lys Arg Met Tyr Ala Gly Leu Lys Asn Ala Gly Glu Thr  
 435 440 445  
 50 Trp Tyr Asp Ile Thr Gly Asn Arg Ser Asp Thr Val Lys Ile Gly Ser  
 450 455 460  
 Asp Gly Trp Gly Glu Phe His Val Asn Asp Gly Ser Val Ser Ile Tyr  
 465 470 475 480  
 55 Val Gln Lys

60 <210> 7  
 <211> 1548  
 <212> DNA  
 <213> Bacillus stearothermophilus  
 65 <220>  
 <221> CDS  
 <222> (1) .. (1548)

<400> 7																	
	gcc	gca	ccg	ttt	aac	ggc	acc	atg	atg	cag	tat	ttt	gaa	tgg	tac	ttg	48
	Ala	Ala	Pro	Phe	Asn	Gly	Thr	Met	Met	Gln	Tyr	Phe	Glu	Trp	Tyr	Leu	
	1				5					10					15		
5	ccg	gat	gat	ggc	acg	tta	tgg	acc	aaa	gtg	gcc	aat	gaa	gcc	aac	aac	96
	Pro	Asp	Asp	Gly	Thr	Leu	Trp	Thr	Lys	Val	Ala	Asn	Glu	Ala	Asn	Asn	
				20					25					30			
10	tta	tcc	agc	ctt	ggc	atc	acc	gct	ctt	tgg	ctg	ccg	ccc	gct	tac	aaa	144
	Leu	Ser	Ser	Leu	Gly	Ile	Thr	Ala	Leu	Trp	Leu	Pro	Pro	Ala	Tyr	Lys	
			35					40					45				
15	gga	aca	agc	cgc	agc	gac	gta	ggg	tac	gga	gta	tac	gac	ttg	tat	gac	192
	Gly	Thr	Ser	Arg	Ser	Asp	Val	Gly	Tyr	Gly	Val	Tyr	Asp	Leu	Tyr	Asp	
		50					55					60					
20	ctc	ggc	gaa	ttc	aat	caa	aaa	ggg	acc	gtc	cgc	aca	aaa	tac	gga	aca	240
	Leu	Gly	Glu	Phe	Asn	Gln	Lys	Gly	Thr	Val	Arg	Thr	Lys	Tyr	Gly	Thr	
	65					70					75					80	
25	aaa	gct	caa	tat	ctt	caa	gcc	att	caa	gcc	gcc	cac	gcc	gct	gga	atg	288
	Lys	Ala	Gln	Tyr	Leu	Gln	Ala	Ile	Gln	Ala	Ala	His	Ala	Ala	Gly	Met	
					85					90					95		
30	caa	gtg	tac	gcc	gat	gtc	gtg	ttc	gac	cat	aaa	ggc	ggc	gct	gac	ggc	336
	Gln	Val	Tyr	Ala	Asp	Val	Val	Phe	Asp	His	Lys	Gly	Gly	Ala	Asp	Gly	
				100					105					110			
35	acg	gaa	tgg	gtg	gac	gcc	gtc	gaa	gtc	aat	ccg	tcc	gac	cgc	aac	caa	384
	Thr	Glu	Trp	Val	Asp	Ala	Val	Glu	Val	Asn	Pro	Ser	Asp	Arg	Asn	Gln	
			115					120					125				
40	gaa	atc	tcg	ggc	acc	tat	caa	atc	caa	gca	tgg	acg	aaa	ttt	gat	ttt	432
	Glu	Ile	Ser	Gly	Thr	Tyr	Gln	Ile	Gln	Ala	Trp	Thr	Lys	Phe	Asp	Phe	
		130					135					140					
45	ccc	ggg	cgg	ggc	aac	acc	tac	tcc	agc	ttt	aag	tgg	cgc	tgg	tac	cat	480
	Pro	Gly	Arg	Gly	Asn	Thr	Tyr	Ser	Ser	Phe	Lys	Trp	Arg	Trp	Tyr	His	
	145					150					155					160	
50	ttt	gac	ggc	gtt	gat	tgg	gac	gaa	agc	cga	aaa	ttg	agc	cgc	att	tac	528
	Phe	Asp	Gly	Val	Asp	Trp	Asp	Glu	Ser	Arg	Lys	Leu	Ser	Arg	Ile	Tyr	
				165						170					175		
55	aaa	ttc	cgc	ggc	atc	ggc	aaa	gcg	tgg	gat	tgg	gaa	gta	gac	acg	gaa	576
	Lys	Phe	Arg	Gly	Ile	Gly	Lys	Ala	Trp	Asp	Trp	Glu	Val	Asp	Thr	Glu	
				180					185					190			
60	aac	gga	aac	tat	gac	tac	tta	atg	tat	gcc	gac	ctt	gat	atg	gat	cat	624
	Asn	Gly	Asn	Tyr	Asp	Tyr	Leu	Met	Tyr	Ala	Asp	Leu	Asp	Met	Asp	His	
			195					200					205				
65	ccc	gaa	gtc	gtg	acc	gag	ctg	aaa	aac	tgg	ggg	aaa	tgg	tat	gtc	aac	672
	Pro	Glu	Val	Val	Thr	Glu	Leu	Lys	Asn	Trp	Gly	Lys	Trp	Tyr	Val	Asn	
		210					215					220					
70	aca	acg	aac	att	gat	ggg	ttc	cgg	ctt	gat	gcc	gtc	aag	cat	att	aag	720
	Thr	Thr	Asn	Ile	Asp	Gly	Phe	Arg	Leu	Asp	Ala	Val	Lys	His	Ile	Lys	
						230					235					240	
75	ttc	agt	ttt	ttt	cct	gat	tgg	ttg	tcg	tat	gtg	cgt	tct	cag	act	ggc	768
	Phe	Ser	Phe	Phe	Pro	Asp	Trp	Leu	Ser	Tyr	Val	Arg	Ser	Gln	Thr	Gly	
					245					250					255		
80	aag	ccg	cta	ttt	acc	gtc	ggg	gaa	tat	tgg	agc	tat	gac	atc	aac	aag	816
	Lys	Pro	Leu	Phe	Thr	Val	Gly	Glu	Tyr	Trp	Ser	Tyr	Asp	Ile	Asn	Lys	

	260	265	270	
5	ttg cac aat tac att acg aaa aca gac gga acg atg tct ttg ttt gat Leu His Asn Tyr Ile Thr Lys Thr Asp Gly Thr Met Ser Leu Phe Asp 275 280 285	864		
10	gcc ccg tta cac aac aaa ttt tat acc gct tcc aaa tca ggg ggc gca Ala Pro Leu His Asn Lys Phe Tyr Thr Ala Ser Lys Ser Gly Gly Ala 290 295 300	912		
15	ttt gat atg cgc acg tta atg acc aat act ctc atg aaa gat caa ccg Phe Asp Met Arg Thr Leu Met Thr Asn Thr Leu Met Lys Asp Gln Pro 305 310 315 320	960		
20	aca ttg gcc gtc acc ttc gtt gat aat cat gac acc gaa ccc ggc caa Thr Leu Ala Val Thr Phe Val Asp Asn His Asp Thr Glu Pro Gly Gln 325 330 335	1008		
25	gcg ctg cag tca tgg gtc gac cca tgg ttc aaa ccg ttg gct tac gcc Ala Leu Gln Ser Trp Val Asp Pro Trp Phe Lys Pro Leu Ala Tyr Ala 340 345 350	1056		
30	ttt att cta act cgg cag gaa gga tac ccg tgc gtc ttt tat ggt gac Phe Ile Leu Thr Arg Gln Glu Gly Tyr Pro Cys Val Phe Tyr Gly Asp 355 360 365	1104		
35	tat tat ggc att cca caa tat aac att cct tcg ctg aaa agc aaa atc Tyr Tyr Gly Ile Pro Gln Tyr Asn Ile Pro Ser Leu Lys Ser Lys Ile 370 375 380	1152		
40	gat ccg ctc ctc atc gcg cgc agg gat tat gct tac gga acg caa cat Asp Pro Leu Leu Ile Ala Arg Arg Asp Tyr Ala Tyr Gly Thr Gln His 385 390 395 400	1200		
45	gat tat ctt gat cac tcc gac atc atc ggg tgg aca agg gaa ggg ggc Asp Tyr Leu Asp His Ser Asp Ile Ile Gly Trp Thr Arg Glu Gly Gly 405 410 415	1248		
50	act gaa aaa cca gga tcc gga ctg gcc gca ctg atc acc gat ggg ccg Thr Glu Lys Pro Gly Ser Gly Leu Ala Ala Leu Ile Thr Asp Gly Pro 420 425 430	1296		
55	gga gga agc aaa tgg atg tac gtt ggc aaa caa cac gct gga aaa gtg Gly Gly Ser Lys Trp Met Tyr Val Gly Lys Gln His Ala Gly Lys Val 435 440 445	1344		
60	ttc tat gac ctt acc ggc aac cgg agt gac acc gtc acc atc aac agt Phe Tyr Asp Leu Thr Gly Asn Arg Ser Asp Thr Val Thr Ile Asn Ser 450 455 460	1392		
65	gat gga tgg ggg gaa ttc aaa gtc aat ggc ggt tcg gtt tcg gtt tgg Asp Gly Trp Gly Glu Phe Lys Val Asn Gly Gly Ser Val Ser Val Trp 465 470 475 480	1440		
70	gtt cct aga aaa acg acc gtt tct acc atc gct cgg ccg atc aca acc Val Pro Arg Lys Thr Thr Val Ser Thr Ile Ala Arg Pro Ile Thr Thr 485 490 495	1488		
75	cga ccg tgg act ggt gaa ttc gtc cgt tgg acc gaa cca cgg ttg gtg Arg Pro Trp Thr Gly Glu Phe Val Arg Trp Thr Glu Pro Arg Leu Val 500 505 510	1536		
80	gca tgg cct tga Ala Trp Pro 515	1548		

15

<210> 8  
 <211> 515  
 <212> PRT  
 <213> *Bacillus stearothermophilus*

5

<400> 8  
 Ala Ala Pro Phe Asn Gly Thr Met Met Gln Tyr Phe Glu Trp Tyr Leu  
 1 5 10 15

10

Pro Asp Asp Gly Thr Leu Trp Thr Lys Val Ala Asn Glu Ala Asn Asn  
 20 25 30

Leu Ser Ser Leu Gly Ile Thr Ala Leu Trp Leu Pro Pro Ala Tyr Lys  
 35 40 45

15

Gly Thr Ser Arg Ser Asp Val Gly Tyr Gly Val Tyr Asp Leu Tyr Asp  
 50 55 60

Leu Gly Glu Phe Asn Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly Thr  
 65 70 75 80

Lys Ala Gln Tyr Leu Gln Ala Ile Gln Ala Ala His Ala Ala Gly Met  
 85 90 95

20

Gln Val Tyr Ala Asp Val Val Phe Asp His Lys Gly Gly Ala Asp Gly  
 100 105 110

Thr Glu Trp Val Asp Ala Val Glu Val Asn Pro Ser Asp Arg Asn Gln  
 115 120 125

30

Glu Ile Ser Gly Thr Tyr Gln Ile Gln Ala Trp Thr Lys Phe Asp Phe  
 130 135 140

Pro Gly Arg Gly Asn Thr Tyr Ser Ser Phe Lys Trp Arg Trp Tyr His  
 145 150 155 160

35

Phe Asp Gly Val Asp Trp Asp Glu Ser Arg Lys Leu Ser Arg Ile Tyr  
 165 170 175

Lys Phe Arg Gly Ile Gly Lys Ala Trp Asp Trp Glu Val Asp Thr Glu  
 180 185 190

Asn Gly Asn Tyr Asp Tyr Leu Met Tyr Ala Asp Leu Asp Met Asp His  
 195 200 205

40

Pro Glu Val Val Thr Glu Leu Lys Asn Trp Gly Lys Trp Tyr Val Asn  
 210 215 220

Thr Thr Asn Ile Asp Gly Phe Arg Leu Asp Ala Val Lys His Ile Lys  
 225 230 235 240

Phe Ser Phe Phe Pro Asp Trp Leu Ser Tyr Val Arg Ser Gln Thr Gly  
 245 250 255

50

Lys Pro Leu Phe Thr Val Gly Glu Tyr Trp Ser Tyr Asp Ile Asn Lys  
 260 265 270

Leu His Asn Tyr Ile Thr Lys Thr Asp Gly Thr Met Ser Leu Phe Asp  
 275 280 285

60

Ala Pro Leu His Asn Lys Phe Tyr Thr Ala Ser Lys Ser Gly Gly Ala  
 290 295 300

Phe Asp Met Arg Thr Leu Met Thr Asn Thr Leu Met Lys Asp Gln Pro  
 305 310 315 320

65

Thr Leu Ala Val Thr Phe Val Asp Asn His Asp Thr Glu Pro Gly Gln

16

325                      330                      335  
 Ala Leu Gln Ser Trp Val Asp Pro Trp Phe Lys Pro Leu Ala Tyr Ala  
                          340                      345                      350  
 5 Phe Ile Leu Thr Arg Gln Glu Gly Tyr Pro Cys Val Phe Tyr Gly Asp  
                          355                      360                      365  
 Tyr Tyr Gly Ile Pro Gln Tyr Asn Ile Pro Ser Leu Lys Ser Lys Ile  
 10                      370                      375                      380  
 Asp Pro Leu Leu Ile Ala Arg Arg Asp Tyr Ala Tyr Gly Thr Gln His  
                          385                      390                      395                      400  
 15 Asp Tyr Leu Asp His Ser Asp Ile Ile Gly Trp Thr Arg Glu Gly Gly  
                          405                      410                      415  
 Thr Glu Lys Pro Gly Ser Gly Leu Ala Ala Leu Ile Thr Asp Gly Pro  
                          420                      425                      430  
 20 Gly Gly Ser Lys Trp Met Tyr Val Gly Lys Gln His Ala Gly Lys Val  
                          435                      440                      445  
 Phe Tyr Asp Leu Thr Gly Asn Arg Ser Asp Thr Val Thr Ile Asn Ser  
 25                      450                      455                      460  
 Asp Gly Trp Gly Glu Phe Lys Val Asn Gly Gly Ser Val Ser Val Trp  
                          465                      470                      475                      480  
 30 Val Pro Arg Lys Thr Thr Val Ser Thr Ile Ala Arg Pro Ile Thr Thr  
                          485                      490                      495  
 Arg Pro Trp Thr Gly Glu Phe Val Arg Trp Thr Glu Pro Arg Leu Val  
                          500                      505                      510  
 35 Ala Trp Pro  
                          515

40 <210> 9  
     <211> 31  
     <212> DNA  
     <213> Artificial Sequence  
 45 <220>  
     <223> Description of Artificial Sequence: Primer

<400> 9  
 50 ggtagtaggc accgtagccc caatccgctt g

31

<210> 10  
 <211> 36  
 55 <212> DNA  
     <213> Artificial Sequence

<220>  
 60 <223> Description of Artificial Sequence: Primer  
     <400> 10  
     ggtagtaggc accgtagccc caatcccatt ggctcg

36

65 <210> 11  
     <211> 28



<212> DNA  
<213> Artificial Sequence

<220>  
5 <223> Description of Artificial Sequence: Primer

<400> 11  
ctgtgactgg tgagtactca accaagtc 28

10 <210> 12  
<211> 31  
<212> DNA  
<213> Artificial Sequence

15 <220>  
<223> Description of Artificial Sequence: Primer

<400> 12  
20 ggtagtaggc accgtagccc tcacccgctt g 31

<210> 13  
<211> 31  
25 <212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

30 <400> 13  
ggtagtaggc accgtagccc ataccgctt g 31

35 <210> 14  
<211> 31  
<212> DNA  
<213> Artificial Sequence

40 <220>  
<223> Description of Artificial Sequence: Primer

<400> 14  
45 ggtagtaggc accgtagcca ataccgctt g 31

<210> 15  
<211> 36  
<212> DNA  
50 <213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

55 <400> 15  
gcagcatgga actgctyatg aagaggcacg tcaaac 36

<210> 16  
60 <211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
65 <223> Description of Artificial Sequence: Primer

<400> 16

catagttgcc gaattcattg gaaacttccc 30

5 <210> 17  
<211> 34  
<212> DNA  
<213> Artificial Sequence

10 <220>  
<223> Description of Artificial Sequence: Primer

<400> 17  
catagttgcc gaattcaggg gaaacttccc aatc 34

15 <210> 18  
<211> 41  
<212> DNA  
<213> Artificial Sequence

20 <220>  
<223> Description of Artificial Sequence: Primer

<400> 18  
25 ccgcgccccg ggaaatcaaa tttgtccag gctttaatta g 41

<210> 19  
<211> 32  
30 <212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

35 <400> 19  
caaaatggta ccaataccac ttaaaatcgc tg 32

40 <210> 20  
<211> 29  
<212> DNA  
<213> Artificial Sequence

45 <220>  
<223> Description of Artificial Sequence: Primer

<400> 20  
50 cttcccaatc ccaagtcttc ccttgaaac 29

<210> 21  
<211> 36  
55 <212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

60 <400> 21  
cttaatttct gctacgacgt caggatggtc ataatc 36

65 <210> 22  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

5 <400> 22  
cgcccaagtc attcgaccag tactcagcta ccgtaaac 38

<210> 23  
10 <211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
15 <223> Description of Artificial Sequence: Primer

<400> 23  
gccgttttca ttgtcgactt cccaatccc 29

20 <210> 24  
<211> 35  
<212> DNA  
<213> Artificial Sequence

25 <220>  
<223> Description of Artificial Sequence: Primer

<400> 24  
30 ggaatttcgc gctgactagt cccgtacata tcccc 35

<210> 25  
<211> 36  
35 <212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

40 <400> 25  
ggcaggaatt tcgcgacctt tcgtcccgta catatc 36

45 <210> 26  
<211> 36  
<212> DNA  
<213> Artificial Sequence

50 <220>  
<223> Description of Artificial Sequence: Primer

<400> 26  
55 cctcattctg cagcagcagc cgtaaattggc acgctg 36

<210> 27  
<211> 38  
60 <212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

65 <400> 27  
ccagacggca gtaataccga tatccgataa atgttccg 38

<210> 28  
<211> 30  
<212> DNA  
5 <213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Primer  
  
10 <400> 28  
cggatatcgg tattactgcc gtctggattc 30  
  
<210> 29  
15 <211> 21  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
20 <223> Description of Artificial Sequence: Primer  
  
<400> 29  
ctcgtcccaa tcggttcggt c 21  
  
25 <210> 30  
<211> 26  
<212> DNA  
<213> Artificial Sequence  
30 <220>  
<223> Description of Artificial Sequence: Primer  
  
<400> 30  
35 gatgtatgcc gacttcgatt atgacc 26  
  
<210> 31  
<211> 30  
40 <212> DNA  
<213> Artificial Sequence  
  
<220>  
<223> Description of Artificial Sequence: Primer  
45 <400> 31  
catagttgcc gaattcattg gaaacttccc 30  
  
50 <210> 32  
<211> 24  
<212> DNA  
<213> Artificial Sequence  
  
55 <220>  
<223> Description of Artificial Sequence: Primer  
  
<400> 32  
60 ccgattgctg acgctgttat ttgc 24  
  
<210> 33  
<211> 25  
<212> DNA  
65 <213> Artificial Sequence  
  
<220>

<223> Description of Artificial Sequence: Primer

<400> 33  
5 gccaaagcgga taacggctac ggtgc 25

<210> 34  
<211> 28  
<212> DNA  
10 <213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

15 <400> 34  
gaacgagcca atcggacgtg ggctacgg 28

<210> 35  
20 <211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
25 <223> Description of Artificial Sequence: Primer

<400> 35  
30 ggaacgagcc aatcggataa cggctacggt gc 32

<210> 36  
<211> 25  
<212> DNA  
<213> Artificial Sequence

35 <220>  
<223> Description of Artificial Sequence: Primer

<400> 36  
40 gcatataagg gactgagcca agcgg 25

<210> 37  
<211> 25  
45 <212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

50 <400> 37  
caaccacaaa gccggcgctg atgcg 25

<210> 38  
<211> 41  
<212> DNA  
<213> Artificial Sequence

55 <220>  
<223> Description of Artificial Sequence: Primer

<400> 38  
60 gcatataagg gactgagcca atcggataac ggctacggtg c 41

65 <210> 39

<211> 28  
<212> DNA  
<213> Artificial Sequence

5 <220>  
<223> Description of Artificial Sequence: Primer

<400> 39  
gaacgagccg atcggacgtg ggctacgg 28

10

<210> 40  
<211> 28  
<212> DNA  
15 <213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

20 <400> 40  
gaacgagcca aaacgacgtg ggctacgg 28

25